

Quantum corrections to neutrino masses and mixing angles¹

Piotr H. Chankowski and Stefan Pokorski

Institute of Theoretical Physics, Warsaw University
Hoża 69, 00-681 Warsaw, Poland

Contents

- 1 Introduction
- 2 Fermion masses and mixing
- 3 Neutrino masses in the effective theory
- 4 Quantum corrections from the renormalization group evolution
 - 4.1 RG equations for the CKM matrix
 - 4.2 RG equations for neutrino masses and mixing angles
 - 4.3 Evolution of the neutrino masses
 - 4.4 Mixing of two neutrinos
 - 4.5 Mixing of three neutrinos and fixed points
- 5 Low energy threshold corrections
 - 5.1 Threshold corrections in the SM
 - 5.2 Threshold corrections in the MSSM
 - 5.2.1 Three-fold degeneracy and flavour diagonal corrections
 - 5.2.2 Three-fold degeneracy and flavour non-diagonal corrections
 - 5.2.3 Two-fold degeneracy and threshold corrections
- 6 Conclusions

Appendix A

Appendix B

References

¹To appear in the review section of Modern Physics A.

1 Introduction

Quark and lepton masses and mixing angles are free parameters of the Standard Model (SM). They are known with various degree of accuracy, from the precision direct measurements of charged lepton masses, a combination of experimental data and theoretical arguments for quark masses and relatively precise measurements of quark mixing angles and the phase parameterizing the violation of CP invariance in the SM. A new element in this picture are neutrino masses and mixing angles. There is at present strong experimental evidence for neutrino oscillations whose most obvious and most natural explanation is that neutrinos have non-zero masses and the neutrino mass eigenstates are different from the weak interaction eigenstates. Although far from having final interpretation, the present experimental data give interesting preliminary information about the neutrino mass sector and the forthcoming experiments are expected to resolve the remaining ambiguities.

On the theoretical side, the origin of the interactions giving rise to fermion masses is a problem that cannot be addressed in the framework of the SM. The physical scale (we shall call it M) of the, still unknown, theory of fermion masses is certainly above the electroweak scale and quite likely it may even be close to the GUT or Planck scales. Such a high scale is suggested, for instance, by the see-saw interpretation of the magnitude of the neutrino masses indicated by experiment. Ultimately, the theory will predict at the scale M the running fermion mass parameters (and perhaps also other parameters) of the effective low energy theory, describing physics at energies $< M$.¹

To relate the mass parameters of the effective low energy theory to the experimentally measured quantities one has to include quantum corrections that already do not depend on the specific theory of fermion masses but only on the low energy effective theory. Clearly, close to the electroweak scale this theory is the Standard Model (SM). One possibility is that the SM remains the correct effective theory up to the scale M . This is conceivable particularly if the scale M is relatively low. Another possibility is that the SM needs to be embedded into a bigger effective theory already much below M . The latter case may, for instance, happen if the low energy supersymmetry is realized in Nature. In this review we discuss quantum corrections to neutrino masses both in the SM and in its supersymmetric extension, the Minimal Supersymmetric Standard Model (MSSM).

Quite generally one can distinguish two classes of quantum corrections that enter on a somewhat different footing. The first one is given by the RGEs describing the evolution

¹The running mass parameters of the effective low energy theory can in principle be calculated in the underlying theory at any renormalization scale Q by proper inclusion of the high energy threshold corrections. Choosing $Q \approx M$ one minimizes those corrections.

of the fermion mass parameters in the SM or MSSM from the scale M down to some scale close to the electroweak scale. Another source of quantum corrections are the so-called low energy threshold effects. Strictly speaking these are the corrections necessary to express measurable quantities like neutrino masses and mixing angles in terms of the running (renormalized) parameters of the effective theory Lagrangian. Formally, their inclusion renders the prediction for observables independent of the choice of the scale to which the RGEs are integrated. In the SM the threshold corrections are unambiguous and, if that scale is taken to be close to M_Z , they are small and can be neglected. Such corrections may, however, be very important in the MSSM since there they depend also on the sparticle masses and couplings.

In Section 2 we review the neutrino masses and mixing and stress the differences with the quark sector. Contrary to the small mixing in the latter one, at least one neutrino mixing angle and, quite likely two, are close to maximal [1]. A maximal or bimaximal mixing would be quite natural for (approximately) degenerate masses, much larger than their differences [2]. Such mass patterns are very different from the hierarchical masses of the charged fermions but are consistent with experiment. Indeed, only mass squared differences can be inferred from the data and not the neutrino masses themselves.

The observed (mixing) and potential (mass pattern) differences between the neutrino and quark sectors provide strong motivation for studying quantum corrections to neutrino masses and mixing. It is well known that the corrections are small for the quarks, just because of their hierarchical masses and mixing. As we shall review in this paper, that remains true for neutrinos if their masses are hierarchical too. However, quantum corrections may give strong, qualitatively new, effects if the neutrino masses are not hierarchical.

The main parts of this review are Sections 4 and 5. In Section 4 we discuss quantum corrections described by the renormalization group evolution and in Section 5 the potential effects of the low energy threshold corrections are reviewed.

A brief overview of quantum corrections in the neutrino sector is given in Section 6.

2 Fermion masses and mixing

For quarks and charged leptons the Particle Data Group [3] gives the following values of the masses²:

$$\begin{aligned} m_u &= 1.5 - 5.0 \text{ MeV}, & m_c &= 1.15 - 1.35 \text{ GeV}, & m_t &= 174.3 \pm 5.1 \text{ GeV}, \\ m_d &= 3.0 - 9.0 \text{ MeV}, & m_s &= 60 - 170 \text{ MeV}, & m_b &= 4.0 - 4.4 \text{ GeV}, \\ m_e &= 511 \text{ keV}, & m_\mu &= 105.7 \text{ MeV}, & m_\tau &= 1.777 \text{ GeV} \end{aligned} \quad (2.1)$$

and of the quark mixing (absolute values of the elements of the Cabibbo-Kobayashi-Maskawa (CKM) matrix):

$$\begin{pmatrix} 0.9742 - 0.9757 & 0.219 - 0.226 & 0.002 - 0.005 \\ 0.219 - 0.225 & 0.9743 - 0.9749 & 0.037 - 0.043 \\ 0.004 - 0.014 & 0.035 - 0.043 & 0.9990 - 0.9993 \end{pmatrix} \quad (2.2)$$

The pattern of quark masses and mixing is clear: hierarchical masses and small (and also) hierarchical mixing: the larger the mass difference, the smaller the mixing angle. In the framework of the SM, fermion masses and mixing angles are used to fix the Yukawa matrices which are the actual free parameters of the theory. They cannot, however, be uniquely reconstructed from the experimental data: there is always the ambiguity of rotating the electroweak basis and all we can have at present is merely a phenomenological parametrization.

Neutrino masses and mixing are inferred from a number of oscillation experiments. The interpretation of neutrino oscillations in terms of massive neutrinos is the most natural one. To discuss this interpretation, let us first recall the oscillation pattern for two hypothetical neutrinos with masses m_1 and m_2 whose quantum fields are linear combinations of two quantum fields ν_A and ν_B that are weak eigenstates

$$\begin{aligned} \nu_1 &= \cos \vartheta \nu_A + \sin \vartheta \nu_B \\ \nu_2 &= -\sin \vartheta \nu_A + \cos \vartheta \nu_B \end{aligned} \quad (2.3)$$

The probabilities that in a given experiment a neutrino produced in the interaction of the charged lepton of flavour A with the $W^\pm$ boson is detected at a distance L as the neutrino creating the charged lepton of flavour B , $B \neq A$, or A are³

$$\begin{aligned} P^{2 \times 2}(\nu_A \rightarrow \nu_B) &= \sin^2 2\vartheta \sin^2 \left(\frac{\Delta m^2 L_{\text{exp}}}{4E_{\text{exp}}} \right) \\ P^{2 \times 2}(\nu_A \rightarrow \nu_A) &= 1 - P^{2 \times 2}(\nu_A \rightarrow \nu_B) \end{aligned} \quad (2.4)$$

²For u , d and s quarks the quoted values are the running masses in the $\overline{\text{MS}}$ scheme at $Q = 2 \text{ GeV}$. The values given for the c and b quarks are the $\overline{\text{MS}}$ running masses at $Q = m_c$ and $Q = m_b$, respectively. Finally m_t given here is the pole mass.

³The formulae like (2.4) are usually derived in the framework of quantum mechanics. The proper

We see that the oscillation probability depends on $\Delta m^2 \equiv m_{\nu_2}^2 - m_{\nu_1}^2$, the distance L_{exp} , the mixing angle ϑ and the neutrino energy E_{exp} . It is clear that for observing the oscillation pattern the factor $\Delta m^2 L/4E$ should be of order $\mathcal{O}(1)$.⁴ Hence, in general, longer oscillation distances are necessary to probe smaller mass squared differences. On the other hand, for $\Delta m^2 L/4E \gg 1$ the transition probability is $P(\nu_A \rightarrow \nu_B) \approx \frac{1}{2} \sin^2 2\vartheta$, i.e. it is insensitive to Δm^2 , because in realistic applications the expression (2.4) has to be averaged over some non-zero interval of the initial neutrino energies ΔE [5].

We can now summarize the results of various experiments. They are usually interpreted in terms of the effective 2×2 parametrization (2.4). Historically, the first information came from the experiments measuring the flux of ν_e neutrinos produced in the Sun (for references see e.g. the review [5]). A convincing evidence for the solar neutrino oscillation was provided by the Kamiokande and Superkamiokande experiments which established a strong ($\approx 50\%$) suppression of the flux of neutrinos from the nuclear reaction ${}^8\text{B} \rightarrow {}^8\text{Be}^* + e^+ + \nu_e$. The most plausible explanation of those results is the transmutation of ν_e 's produced in the core of the Sun into another type of neutrinos such as ν_μ , ν_τ and/or the so-called sterile neutrino ν_{sterile} which does not interact with the $W^\pm$ or Z^0 bosons. Such a transmutation can occur either during their flight from the Sun to the Earth (the so-called vacuum oscillations (VO)) or through the resonant transition in the matter of the outer layers of the Sun (the so-called MSW effect [6]). As follows from the formulae (2.4) with $A \equiv e$ and $B \equiv \mu, \tau$ or s , in the case of VO with $L \approx 1.5 \times 10^8$ km (the Sun-Earth mean distance) and for mean ${}^8\text{B}$ neutrino energy of order $E \sim 10$ MeV, the solar neutrino experiments are sensitive to $\Delta m_{\text{sol}}^2 \sim \mathcal{O}(10^{-(11-10)})$ eV². The deficit is then explained with $\sin^2 2\vartheta_{\text{sol}} > 0.7$. In the case of the MSW resonant conversion the formulae (2.4) for the transition and survival probabilities are replaced by more complicated expressions [5] which depend also on the electron and neutron number densities in the Sun. The observed ν_e neutrino deficit can be then explained either for $\Delta m_{\text{sol}}^2 \sim \mathcal{O}(10^{-5})$ eV² and $\sin^2 2\vartheta_{\text{sol}} \sim \mathcal{O}(10^{-(3-2)})$ (the so-called SAMWS solution) or for $\Delta m_{\text{sol}}^2 \sim \mathcal{O}(10^{-4})$ eV² and $\sin^2 2\vartheta_{\text{sol}} > 0.5$ (the so-called LAMWS solution).

picture of neutrino oscillations is however the field theoretical one [4]. Due to the field mixing (2.3) there is in general a nonzero amplitude for emission (absorption) of any of the neutrino mass eigenstates ν_a in the interaction of charged lepton of flavour A with the W -boson. The change of the neutrino flavour (inferred from the flavours of the charged leptons) is in this picture due to the coherent sum of Feynman diagrams describing exchanges of all virtual mass eigenstates of neutrinos between the emission and detection vertices. Taking properly into account the effects of wave packets describing initial and final states it can be shown that the formula (2.4) is in most cases a sufficient approximation to the full result, which automatically accounts for decoherence effects and depends on the overlap of the wave packets describing the initial and final states [4].

⁴Recall that $\frac{\Delta m^2 L}{4E} = 1.27 \times \frac{(\Delta m^2/1 \text{ eV}^2)(L/1 \text{ km})}{(E/1 \text{ GeV})}$.

The deficit of neutrinos was also revealed by the measurements of the flux of ν_μ and $\bar{\nu}_\mu$ neutrinos produced together with ν_e 's in the Earth's atmosphere by the cosmic rays. The results of the Superkamiokande experiment [7] are most easily explained by ν_μ oscillation into another type of neutrinos. The oscillatory explanation is further supported by the zenith angle dependence of the ν_μ and $\bar{\nu}_\mu$ flux deficit. For typical ν_μ energies of order $\sim \text{GeV}$ and $20 \text{ km} \lesssim L \lesssim 1.3 \times 10^4 \text{ km}$, the observed ν_μ deficit is explained for $\Delta M_{\text{atm}}^2 \approx 3.2 \times 10^{-3} \text{ eV}^2$ and $\sin^2 2\vartheta_{\text{atm}} > 0.82$. Also, the Superkamiokande data seem to favour $\nu_\mu \rightarrow \nu_\tau$ oscillations over $\nu_\mu \rightarrow \nu_{\text{sterile}}$ [8].

Neutrino oscillations are also intensively searched for in various reactor or accelerator based experiments (for review see [5, 9]). Except for the LSND experiment reporting [10] a positive signal for the $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations, which would require $0.1 \text{ eV}^2 \lesssim \Delta M^2 \lesssim 1 \text{ eV}^2$ for the oscillatory explanation, the results of all other experiments are negative. Particularly strong constraint comes from the CHOOZ reactor experiment. It excludes disappearance of $\bar{\nu}_e$ with mean energies $\sim 3 \text{ MeV}$ at the distance $L \approx 1 \text{ km}$. In the 2×2 oscillation framework this translates into the limit

$$\sin^2 2\vartheta_{\text{react}} < 0.1 \quad \text{for} \quad \Delta M^2 \gtrsim 10^{-3} \text{ eV}^2 \quad (2.5)$$

Except for the unconfirmed LSND result, all experiments are consistent with oscillations between the three known neutrino flavours. Moreover, recent SNO measurement [11] of the total solar neutrino flux (i.e. the flux of ν_e , ν_μ and ν_τ) combined with the Superkamiokande data [12] strongly supports this assumption. Therefore, in the rest of the review we assume that the neutrino sector of the low energy effective theory consists of 3 active neutrinos only.

Within the 3×3 framework the solar, atmospheric and reactor data become interrelated and one can draw more definite conclusions. To make the discussion of the neutrino experiments more complete we recall the formula for the transition probabilities $P(\nu_A \rightarrow \nu_B)$ for the case of 3 neutrinos:

$$\begin{aligned} P(\nu_A \rightarrow \nu_B, L, E) &= \left| \sum_a U_{Ba} e^{-i \frac{m_{\nu_a}^2 L}{2E}} U_{Aa}^* \right|^2 \\ &= \sum_a |U_{Ba} U_{Aa}^*|^2 + \text{Re} \sum_a \sum_{b \neq a} U_{Ba} U_{Bb}^* U_{Aa}^* U_{Ab} e^{i \Delta m_{ab}^2 L / 4E} \end{aligned} \quad (2.6)$$

where $\Delta m_{ab}^2 \equiv m_{\nu_a}^2 - m_{\nu_b}^2$. The complex matrix elements are defined in the basis in which charged lepton mass matrix is diagonal, by the decomposition of the ν_A neutrino fields belonging to $SU_L(2)$ doublets into the mass eigenstates field:

$$\nu_A = \sum_a U_{Aa} \nu_a. \quad (2.7)$$

For a real matrix U the expression (2.6) can be written as

$$P(\nu_A \rightarrow \nu_B, L, E) = \delta_{AB} - 4 \sum_{a>b=1}^3 U_{Aa} U_{Ba} U_{Ab} U_{Bb} \sin^2 \left(\frac{\Delta m_{ab}^2 L}{4E} \right). \quad (2.8)$$

To simplify the notation we will denote: $\Delta m^2 \equiv \Delta m_{\text{sol}}^2 \equiv |m_2^2 - m_1^2|$, $\Delta M^2 \equiv \Delta M_{\text{atm}}^2 \equiv |m_3^2 - m_2^2|$. For $\Delta m^2 \ll \Delta M^2$ the measurements of the experiments listed above can now be summarized as follows:

$$\begin{aligned} P_{\text{sol}}(\nu_e \rightarrow \nu_e) &= 1 - 2 \left(1 - U_{13}^2 \right) U_{13}^2 - 4 U_{11}^2 U_{12}^2 \sin^2 \left(\frac{\Delta m^2 L_{\text{sol}}}{4E_{\text{sol}}} \right) \\ P_{\text{atm}}(\nu_\mu \rightarrow \nu_\mu) &= 1 - 4 \left(1 - U_{23}^2 \right) U_{23}^2 \sin^2 \left(\frac{\Delta M^2 L_{\text{atm}}}{4E_{\text{atm}}} \right) \\ P_{\text{react}}(\nu_e \rightarrow \nu_e) &= 1 - 4 \left(1 - U_{13}^2 \right) U_{13}^2 \sin^2 \left(\frac{\Delta M^2 L_{\text{react}}}{4E_{\text{react}}} \right) \end{aligned} \quad (2.9)$$

A convenient parametrization for the mixing matrix U is [14] (see also the next Section):

$$U_{\text{MNS}} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \times \Phi \quad (2.10)$$

where $c_{ij} \equiv \cos \theta_{ij}$ ($s_{ij} \equiv \sin \theta_{ij}$) and Φ is a diagonal matrix $\Phi = \text{diag}(e^{i\alpha_1}, e^{i\alpha_2}, 1)$ (or a similar diagonal matrix with two phases and $\Phi_{11} = 1$ or $\Phi_{22} = 1$). Our convention is such that for vanishing mixing angles θ_{ij} we have $\nu_1 = \nu_e$, $\nu_2 = \nu_\mu$ and $\nu_3 = \nu_\tau$, i.e. no ordering of the neutrino masses m_{ν_a} is assumed. CP is conserved if $\delta = 0 \bmod \pi$ and $\alpha_{1,2} = 0 \bmod \pi/2$. The relation to the effective 2×2 parametrization then reads

$$\begin{aligned} \sin^2 2\vartheta_{\text{react}} &= \sin^2 2\theta_{13} \\ \sin^2 2\vartheta_{\text{atm}} &= \sin^2 2\theta_{23} \left(\cos^2 \theta_{13} + \frac{\sin^2 2\theta_{13}}{4 \cos^2_{23}} \right) \\ \sin^2 2\vartheta_{\text{sol}} &\approx \sin^2 2\theta_{12} \left(1 - 2 \sin^2 \theta_{13} \right) \end{aligned} \quad (2.11)$$

In the limit of small θ_{13} angle imposed by the CHOOZ result (2.5) (non-negligible entries U_{23} and U_{33} and unitarity of U exclude the possibility of $|\theta_{13}| \approx \pi/2$) we get

$$\begin{aligned} \sin^2 2\vartheta_{\text{atm}} &\approx \sin^2 2\theta_{23} \\ \sin^2 2\vartheta_{\text{sol}} &\approx \sin^2 2\theta_{12} \end{aligned} \quad (2.12)$$

Finally we mention that the neutrino Majorana mass term of the form $\mathcal{L}_{\text{mass}} = -\frac{1}{2} m_\nu^{AB} \nu_A \nu_B + \text{H.c}$ violates the lepton number conservation and leads to the neutrino-less double beta decay. Non-observation of such decays implies [15]

$$|m_\nu^{ee}| < 0.35(0.27) \text{ eV} \quad \text{at } 90\% \text{ (68\%)} \text{ C.L.}, \quad (2.13)$$

i.e.

$$|m_{\nu_1} c_{12}^2 c_{13}^2 e^{2i\alpha_1} + m_{\nu_2} s_{12}^2 c_{13}^2 e^{2i\alpha_2} + m_{\nu_3} s_{13}^2 e^{2i\delta}| < 0.35(0.27) \text{ eV at } 90\% \text{ (68\%)} \text{ C.L.} \quad (2.14)$$

Concluding this section, we stress that the pattern of neutrino mixing is distinctly different from the one observed in the quark sector. At least one mixing angle (θ_{23}) is large and the recent data favour solutions with two large mixing angles [1]. The data indicate $\Delta m^2 \ll \Delta M^2$ but, since the masses are not measured, several options for the values of the masses themselves are still possible:

- i) $\Delta M^2 \approx m_{\nu_3}^2 \gg m_{\nu_{2(1)}}^2 \gg m_{\nu_{1(2)}}^2$ (hierarchical)
- ii) $\Delta M^2 \approx m_{\nu_1}^2, m_{\nu_2}^2 \gg m_{\nu_3}^2$ or $\Delta M^2 \approx m_{\nu_3}^2 \gg m_{\nu_1}^2, m_{\nu_2}^2$, with $m_{\nu_1}^2, m_{\nu_2}^2 \gg \Delta m^2$ (partly degenerate)
- iii) $m_{\nu_3}^2 \approx m_{\nu_2}^2 \approx m_{\nu_1}^2$ and all of order or larger than ΔM^2 (degenerate).

Special cases of the partly degenerate and degenerate patterns are equal masses, $|m_{\nu_1}| = |m_{\nu_2}|$ or $|m_{\nu_1}| = |m_{\nu_2}| = |m_{\nu_3}|$, respectively. In our language we refer to them as two-fold and three-fold degeneracies, to distinguish them from, more general, partly degenerate and degenerate pattern defined by *ii*) and *iii*). The last two possibilities are very different from the pattern known from the quark sector but almost maximal (bimaximal) mixing suggested by the experimental data makes them an interesting alternative to the hierarchical pattern. It is important to remember that in that respect neutrinos may be qualitatively different from quarks because, unlike the charged fermions, they can be Majorana particles. As we shall see, the magnitude and the importance of quantum corrections depends strongly on the assumed pattern of the masses.

3 Neutrino masses in the effective theory

We begin by recalling how the quark masses are incorporated in the SM (or the MSSM) Lagrangian. We shall use the Weyl spinor notation (see e.g. [16]), which is particularly convenient for Majorana particles. The relation to the standard Dirac notation is explained in the Appendix A. The original Lorentz and $SU(3) \times SU(2) \times U(1)$ invariant Lagrangian contains the Yukawa interactions of the form

$$\mathcal{L}_{\text{Yuk}} = -\epsilon_{ij} H_i u^{cA} \mathbf{Y}_u^{AB} q_j^B - H_j^* d^{cA} \mathbf{Y}_d^{AB} q_j^B - H_j^* e^{cA} \mathbf{Y}_e^{AB} l_j^B + \text{H.c.} \quad (3.1)$$

where $\mathbf{Y}_u$, $\mathbf{Y}_d$, $\mathbf{Y}_e$ are a priori arbitrary complex Yukawa matrices, $\epsilon_{12} = -\epsilon_{21} = -1$, the upper case letters enumerate the three generations of the matter fermions

$$u^c, \quad d^c, \quad e^c, \quad q \equiv \begin{pmatrix} u \\ d \end{pmatrix} \quad \text{and} \quad l \equiv \begin{pmatrix} \nu \\ e \end{pmatrix} \quad (3.2)$$

described by the left-handed Weyl spinors and H is the SM Higgs doublet. In the MSSM there are two Higgs doublets, $H^{(u)}$ and $H^{(d)}$, and one has to replace: $H_i \rightarrow H_i^{(u)}$ and $H_j^* \rightarrow \epsilon_{ij} H_i^{(d)}$. Replacing the Higgs doublet H (doublets $H^{(u)}$ and $H^{(d)}$ in the MSSM) by its vacuum expectation value leads to the following mass terms in the SM (MSSM) Lagrangian:

$$\mathcal{L}_{\text{mass}} = -u^{cA}(v\mathbf{Y}_u)^{AB}u^B - d^{cA}(v\mathbf{Y}_d)^{AB}d^B - e^{cA}(v\mathbf{Y}_e)^{AB}e^B + \text{H.c.} \quad (3.3)$$

(In the MSSM $v\mathbf{Y}_u \rightarrow v_u\mathbf{Y}_u$, $v\mathbf{Y}_{d,e} \rightarrow -v_d\mathbf{Y}_{d,e}$.) The fermion mass matrices are diagonalized by the unitary chiral rotations

$$\begin{aligned} u &\rightarrow \mathbf{U}_L u, & u^c &\rightarrow u^c \mathbf{U}_R^\dagger, \\ d &\rightarrow \mathbf{D}_L d, & d^c &\rightarrow d^c \mathbf{D}_R^\dagger, \\ e &\rightarrow \mathbf{E}_L e, & e^c &\rightarrow e^c \mathbf{E}_R^\dagger, \end{aligned} \quad (3.4)$$

which give:

$$\begin{aligned} \mathbf{U}_R^\dagger \mathbf{Y}_u \mathbf{U}_L &= \text{diag}(y_u, y_c, y_t) \\ \mathbf{D}_R^\dagger \mathbf{Y}_d \mathbf{D}_L &= \text{diag}(y_d, y_s, y_b) \\ \mathbf{E}_R^\dagger \mathbf{Y}_e \mathbf{E}_L &= \text{diag}(y_e, y_\mu, y_\tau). \end{aligned} \quad (3.5)$$

The only remnant of the nontrivial Yukawa matrices is then the CKM matrix

$$V_{\text{CKM}} \equiv V \equiv \mathbf{U}_L^\dagger \mathbf{D}_L \quad (3.6)$$

appearing in the interactions of the charged quark currents with (massive) charged vector bosons $W_\mu^\pm$. It can be shown (see e.g. [16]) that the CKM matrix can be parameterized by 3 angles and one phase. The other $6 - 1 = 5$ phases can be absorbed in the redefinition of the quark fields.

In the SM defined as a renormalizable theory, with the neutrino and Higgs fields transforming as components of doublets of the $SU_L(2)$ gauge symmetry, neutrinos remain massless. Moreover, as a consequence of the gauge symmetry, renormalizability and the field content, the Lagrangian has two accidental global $U(1)$ symmetries that ensure baryon and lepton number conservation. In the absence of any mass terms for the neutrino fields ν^A , one can always perform the rotation $\nu \rightarrow \mathbf{V}_L \nu$ with $\mathbf{V}_L = \mathbf{E}_L$ so that the counterpart of the CKM matrix in the leptonic sector is trivial.

There are two easy possibilities for extending the particle content of the SM so that neutrino mass is generated by renormalizable interactions. One is to couple two lepton $SU_L(2)$ doublets to a $SU_L(2)$ Higgs triplet (singlets would violate the electric charge

conservation). This possibility is not particularly attractive for several reasons. The smallness of the neutrino masses would require either $v_{\text{triplet}} \ll v_{\text{doublet}}$ or the triplet couplings orders of magnitude smaller than the other Yukawa couplings. Moreover, the introduction of the triplet would make the parameter ρ a free parameter of the theory (hence not calculable) with its own counterterm allowing to adjust its value at will. The other possibility is to introduce a number of new $SU_L(2) \times U_Y(1)$ singlet left-handed leptonic fields ν^c to the Lagrangian and couple them to the Higgs doublet in the same way as the quark fields u^c :

$$\Delta\mathcal{L}_{\text{Yuk}} = -\epsilon_{ij} H_i \nu^{cK} \mathbf{Y}_\nu^{KA} l_j^A + \text{H.c.} \quad (3.7)$$

where $K = 1, \dots, K_{\text{max}}$ and in principle K_{max} could be arbitrary. The interaction (3.7) preserves the known structure of the SM and, if $K = 1, 2, 3$, the symmetry between quarks and leptons is restored. For singlet fields ν^c the Majorana mass term

$$\Delta\mathcal{L}_{\text{Maj}} = -\frac{1}{2} \mathbf{M}_{\text{Maj}}^{KL} \nu^{cK} \nu^{cL} + \text{H.c.} \quad (3.8)$$

can also be added to the Lagrangian. In general both terms should be included in the theory defined by the $SU_L(2) \times U_Y(1)$ gauge symmetry and renormalizability. The Majorana mass term can be, however, eliminated by imposing the additional global $U(1)$ symmetry ensuring conservation of the lepton number L . The fields ν^c must then have $L = -1$, opposite to L of the leptonic doublets l_i . Their complex conjugate $\bar{\nu}^c$ (see Appendix A) can be then interpreted as the right-handed neutrinos. Neutrinos are then Dirac particles like the other fermions. There are two reasons why Dirac masses are not so attractive. One is, again, the need for very small numerical values of the Yukawa couplings $\mathbf{Y}_\nu$. The other is that the lepton number conservation has to be imposed as an additional global symmetry (remember that in the SM it is a consequence of the field content and of the renormalizability and not an additional assumption).

The presence of the Majorana mass term, which breaks the global $U(1)$ symmetry, inevitably makes neutrinos Majorana particles (in fact what allows to interpret in the SM the two helicity states described by the ν field as a particle and an anti-particle is just the lepton number!) Taking for simplicity three⁵ singlet neutrinos ν^c and replacing H ($H^{(u)}$ in the MSSM) by its vacuum expectation value, the general form of the neutrino mass matrix (in the absence of Higgs triplets) is

$$\mathcal{M}_\nu = \begin{pmatrix} 0 & m_D \\ m_D^T & M_{\text{Maj}} \end{pmatrix} \quad (3.9)$$

⁵The formula (3.9) is easily generalized to an arbitrary number of singlet neutrinos ν^c .

where $m_D \equiv v \mathbf{Y}_\nu$ and M_{Maj} are 3×3 matrices in the basis (ν, ν^c) . If both these matrices are diagonal with eigenvalues m_A and M_A , respectively, the physical neutrinos are mixtures of ν_A and ν_A^c with the masses

$$m_{\nu^A \pm} = \frac{1}{2} \left(M_A \pm \sqrt{M_A^2 + 4m_A^2} \right). \quad (3.10)$$

In general, each of the six leptonic fields ν_A and ν_K^c is a linear combination of the six Majorana neutrinos. There is the possibility of oscillations $\nu \rightarrow \nu^c$. The ν^c neutrinos are singlets of the $SU_L(2) \times U(1)$ gauge group and are sterile neutrinos.⁶ The situation simplifies for $M \gg v$ where M is the overall scale of the M_{Maj} entries. For simplicity, we identify it with the introduced earlier scale M , although they could actually be different scales. In this limit there are two sets of mass eigenvalues. They are of the order $\mathcal{O}(M)$ and $\mathcal{O}(v^2/M)$, respectively, and three of the mass eigenstates are very heavy and decouple from the physics at the electroweak scale. This is the so-called see-saw mechanism [17]
⁷ The oscillations occur then effectively among the three “active” neutrinos ν (the ν^c ’s composition is dominated by heavy eigenstates).

It is instructive to repeat the decoupling procedure in the field theoretical language. The effects of the Born diagram, Fig. 1, present in the full theory valid above the scale M are, up to $\mathcal{O}(1/M^2)$, reproduced in the effective theory, describing physics below the scale M , by a non-renormalizable operator of dimension 5 [19]:

$$\Delta \mathcal{L} = -\frac{1}{4M} \mathbf{C}^{AB} (\epsilon_{ij} H_i l_j^A) (\epsilon_{lk} H_l l_k^B) + \text{H.c.} \quad (3.11)$$

in which

$$M^{-1} \mathbf{C} = \mathbf{Y}_\nu^T (\mathbf{M}_{\text{Maj}})^{-1} \mathbf{Y}_\nu. \quad (3.12)$$

The overall scale M of the entries of the matrix $\mathbf{M}_{\text{Maj}}$ has been factorized out from $\mathbf{C}$ to make it dimensionless. Thus, if the scale M is high enough, the effects of the Majorana masses can be discussed in the SM or MSSM supplemented by the operator (3.11). We should also stress that this operator is the only one of dimension 5 contributing to the neutrino Majorana mass matrix. Other possible contributions are of higher dimension [20]. Thus, one may expect that in the effective SM or MSSM the neutrino masses are indeed described by the operator (3.11), even if its origin is different from the see-saw

⁶Note that such oscillations exist only for Majorana neutrinos and are forbidden by L conservation for Dirac neutrinos i.e. the right-handed neutrinos $\bar{\nu}^c$ have correct L but wrong chirality to oscillate into ν .

⁷However in some cases, one of the formally $\mathcal{O}(M)$ mass eigenvalues can remain small as a result of some symmetries of the underlying theory of neutrino masses. The corresponding light state can play the role of a sterile neutrino [18].

mechanism. In particular, degenerate or partly degenerate patterns of neutrino masses require some interplay between the parameters entering the formula (3.12), but one can also imagine that such patterns originate from other mechanisms at the scale M .




Figure 1: Diagram generating the dimension 5 operator (3.11).

We end this section with the discussion of the parametrization of the neutrino mixing matrix. In the effective theory, after the electroweak symmetry breaking, the operator (3.11) is the source of the Majorana mass for the three active neutrinos ν , $\mathcal{L}_{\text{mass}} = -\frac{1}{2} \frac{v^2}{4M} \mathbf{C}^{AB} \nu^A \nu^B + \text{H.c.}$ This mass matrix can be diagonalized by an additional unitary rotation $\nu^A \rightarrow U^{Aa} \nu_a$. Recall we work in the basis in which the leptonic Yukawa coupling is already diagonal and the matrix $\mathbf{C}$ is assumed to be given in the same basis. We have:

$$\frac{v^2}{4M} (U^T \mathbf{C} U)^{ab} = \frac{v^2}{4M} C^a \delta^{ab} = m_{\nu_a} \delta^{ab} \quad (3.13)$$

The matrix U will therefore appear in the couplings of neutrinos to $W^\pm$ bosons and charged Goldstone (and Higgs) boson.⁸ Being unitary, the matrix U depends on 3 angles and 6 phases and can be conveniently written as $U^{Aa} = e^{i\varphi_A} U_{\text{MNS}}^{Aa}$ where U_{MNS} is given in eq. (2.10). Contrary to the quark case, if the Majorana masses m_{ν_a} in eq. (3.13) are to be real and positive, the only freedom that remains is the possibility to re-phase independently the three Dirac fields of the charged leptons $\psi_{e_A} \rightarrow \exp(i\varphi_A) \psi_{e_A}$ (i.e. $e_A \rightarrow \exp(i\varphi_A) e_A$ for the left-handed and $e_A^c \rightarrow \exp(-i\varphi_A) e_A^c$ for their right-handed components). It is then the MNS matrix (2.10) with three angles and three phases that enters the interactions of neutrinos with the SM particles. In the MSSM, however, the phases φ_A can be eliminated (by appropriate rotations of the superpartner fields) only if the slepton mass matrices and charged lepton mass matrices are simultaneously diagonal. In the general case, the phases φ_A appear in the neutrino-sneutrino-neutralino and neutrino-chargino-charged slepton vertices.

⁸In a general electroweak basis in which the leptonic Yukawa coupling is not necessarily diagonal the matrix U is given by the product $U = \mathbf{E}_L^\dagger \mathbf{V}_L$ where the rotations $e \rightarrow \mathbf{E}_L e$ and $\nu \rightarrow \mathbf{V}_L \nu$ diagonalize the mass matrices of charged leptons and neutrinos, respectively.

It is sometimes convenient to work with complex neutrino masses m_{ν_a} in eq. (3.13). The phases $\alpha_{1,2}$ from eq. (2.10) are then absorbed into the masses: $m_{\nu_{1,2}} = |m_{\nu_{1,2}}| \exp(-2i\alpha_{1,2})$. In particular, when CP is conserved the phases $\alpha_{1,2} = 0, \pm\pi/2$ reflecting different CP parities of different neutrinos, can be absorbed into the mass eigenvalues and make them real positive or negative. The MNS matrix is then real. Since only relative signs of the masses play a role, we will for definiteness fix our convention so that m_{ν_3} is always positive. Such a parametrization simplifies greatly the qualitative analysis of the renormalization group equations discussed in the next section.

The value of the mass M is not known. One can expect that it is determined by a beyond the SM theory that provides the physical cut-off for the SM. The idea of Grand Unification and of the big desert would suggest very high value of M . However there are also other models (large extra dimensions) in which M is $\mathcal{O}(1 \text{ TeV})$. In this review we assume that whatever the theory of neutrino masses is, the scale M is high enough to justify the description of the neutrino mass effects by the effective theory (SM or MSSM) with the single operator (3.11) added.

Finally, one should also mention that in the MSSM neutrino mass can originate from R-parity violating interactions [21]. Such models are usually based on low energy mechanisms for neutrino mass generation and will not be discussed in this review.

4 Quantum corrections from the renormalization group evolution

The neutrino mass parameters given at some scale M in the effective theory (SM or MSSM) supplemented by the operator (3.11), determine the measurable quantities after inclusion of quantum corrections. The first category of corrections are those described by the evolution of the effective theory parameters from the scale M down to some low energy scale close to the electroweak scale. They depend on arbitrary powers of the large logarithms of the ratio M/M_Z and are resummed to all orders of the perturbation expansion by means of the renormalization group equations (RGE). The second class of corrections are the so-called low energy threshold corrections. In the SM they connect the mass parameters of the SM Lagrangian renormalized at the scale M_Z to the fermion masses in the effective theory obtained after the electroweak symmetry breaking (whose renormalizable part is QCD+QED) renormalized at the same scale M_Z . In the MSSM, threshold corrections include also the superpartner contributions due to their mass splittings and/or flavour violation in slepton mass matrices. The relative magnitude of the two types of corrections will be discussed later on. In this section we assume that the RG corrections are

the dominant ones and, after deriving the relevant RG equations, we discuss the potential role played by these corrections for the neutrino masses and mixing.

4.1 RG equations for the CKM matrix

Before we derive the RGE for the neutrino masses and mixing we discuss the technically easier case of the evolution of the Yukawa matrices and the CKM matrix.

The renormalization group equations for the Yukawa couplings of the SM and the MSSM at one [22, 23] and two loop [23, 24] are well known. For convenience we reproduce the one loop RGE in Appendix B. They have a matrix structure which encodes the evolution of their eigenvalues and of all entries of the CKM mixing matrix. From the full RGE in the matrix form it is possible [25, 26] to derive the RGE for the eigenvalues of the Yukawa matrices and for the CKM matrix. It is often very instructive to discuss the evolution of those Lagrangian parameters that can be directly determined from the data. For instance, the experimentally known information can be unambiguously extrapolated to any high scale. Moreover, certain qualitative features of the Yukawa coupling pattern necessary to reproduce the experimental data can be easier to understand. The derivation goes as follows. One writes down auxiliary RGEs for the matrices $\mathbf{U}_{L,R}$, $\mathbf{D}_{L,R}$ defined in eq. (3.4) in the form:

$$\begin{aligned}\frac{d}{dt}\mathbf{U}_L &= \mathbf{U}_L \varepsilon_L^U, & \frac{d}{dt}\mathbf{U}_R &= \mathbf{U}_R \varepsilon_R^U, \\ \frac{d}{dt}\mathbf{D}_L &= \mathbf{D}_L \varepsilon_L^D, & \frac{d}{dt}\mathbf{D}_R &= \mathbf{D}_R \varepsilon_R^D,\end{aligned}\tag{4.1}$$

where the matrices $\varepsilon_{L,R}^{U,D}$ are antihermitean, in order to preserve unitarity of the matrices $\mathbf{D}_{L,R}$, and $\mathbf{U}_{L,R}$, and $t = (1/16\pi^2) \ln(Q/M_Z)$. One then looks for the evolution of the $\mathbf{U}_{L,R}$ and $\mathbf{D}_{L,R}$ matrices such that Yukawa matrices remain diagonal during the evolution. Differentiating the four relations (obtained from eqs. (3.5))

$$\begin{aligned}\text{diag}(y_u^2, y_c^2, y_t^2) &= \mathbf{U}_L^\dagger \mathbf{Y}_u^\dagger \mathbf{Y}_u \mathbf{U}_L = \mathbf{U}_R^\dagger \mathbf{Y}_u \mathbf{Y}_u^\dagger \mathbf{U}_R \\ \text{diag}(y_d^2, y_s^2, y_b^2) &= \mathbf{D}_L^\dagger \mathbf{Y}_d^\dagger \mathbf{Y}_d \mathbf{D}_L = \mathbf{D}_R^\dagger \mathbf{Y}_d \mathbf{Y}_d^\dagger \mathbf{D}_R\end{aligned}\tag{4.2}$$

and requiring the derivatives of the diagonal matrices on the rhs to be also diagonal matrices one gets the matrices $\varepsilon_L^{U,D}$:

$$\begin{aligned}(\varepsilon_L^U)_{JI} &= -u_d \frac{y_{u_J}^2 + y_{u_I}^2}{y_{u_J}^2 - y_{u_I}^2} \sum_K V^{JK} V^{IK*} y_{d_K}^2, & (\varepsilon_L^U)_{JJ} &= 0 \\ (\varepsilon_L^D)_{KL} &= -d_u \frac{y_{d_K}^2 + y_{d_L}^2}{y_{d_K}^2 - y_{d_L}^2} \sum_J y_{u_J}^2 V^{JK*} V^{JL}, & (\varepsilon_L^D)_{KK} &= 0\end{aligned}\tag{4.3}$$

and similarly the matrices $\varepsilon_R^{U,D}$:

$$\begin{aligned} (\varepsilon_R^U)_{JI} &= -u_d \frac{2y_{u_J}y_{u_I}}{y_{u_J}^2 - y_{u_I}^2} \sum_K V^{JK} V^{IK*} y_{d_K}^2, & (\varepsilon_R^U)_{JJ} &= 0 \\ (\varepsilon_R^D)_{KL} &= -d_u \frac{2y_{d_K}y_{d_L}}{y_{d_K}^2 - y_{d_L}^2} \sum_J y_{u_J}^2 V^{JK*} V^{JL}, & (\varepsilon_R^D)_{KK} &= 0 \end{aligned} \quad (4.4)$$

where $u_d = d_u = -3/2$ for the SM and $u_d = d_u = 1$ for the MSSM. The evolution of the CKM mixing matrix V is then given by the simple matrix equation

$$\frac{d}{dt}V = -\varepsilon_L^U V + V \varepsilon_L^D. \quad (4.5)$$

Because of the hierarchical pattern of quark masses the expressions (4.3) for $\varepsilon_L^{U,D}$ can be simplified to

$$\begin{aligned} (\varepsilon_L^D)_{KL} &= -d_u y_t^2 V^{3K*} V^{3L} \quad \text{for } K > L & \text{and } (\varepsilon_L^D)_{KL} &= -(\varepsilon_L^D)_{LK} \quad \text{for } K < L \\ (\varepsilon_L^U)_{JI} &= -u_d V^{J3} V^{I3*} y_b^2 \quad \text{for } J > I & \text{and } (\varepsilon_L^U)_{JI} &= -(\varepsilon_L^U)_{IJ} \quad \text{for } J < I. \end{aligned} \quad (4.6)$$

Taking next into account the hierarchy of the entries of the CKM matrix it is easy to find that, to a good approximation, the 2×2 submatrix describing the mixing of the first two generations as well as the element V^{33} do not evolve, while the evolution of the remaining entries is universal and is given by [26, 27]:

$$V(t) = V(0) \exp \left\{ - \int_0^t \left[u_d y_b^2(t') + d_u y_t^2(t') \right] dt' \right\} \quad (4.7)$$

where $V = V^{31}, V^{32}, V^{13}$ or V^{32} . It follows that to one loop the Jarlskog invariant $J \equiv \text{Im}(V^{ud}V^{cs}V^{us*}V^{cd*})$ does not change. By using e.g. the standard parametrization of the CKM matrix it is also easy to derive the RGE for the three mixing angles and the CP violating phase.⁹ For the SM and the MSSM this has been done up to two-loops in ref. [27].

4.2 RGEs for neutrino masses and mixing

The Majorana mass term for neutrinos arises from the dimension 5 operator (3.11). The RGE for $\mathbf{C}^{AB}$ in the SM has been correctly computed only recently in ref. [29] (previous

⁹In the one loop approximation the running of the CKM matrix elements looks particularly simple in the Wolfenstein parametrization [28] (see e.g. [16]) in which only the parameter A changes with the scale.

calculations [30, 31, 32] have errors) and in the MSSM in refs. [31, 32].¹⁰

$$\frac{d}{dt}\mathbf{C} = -K\mathbf{C} - \kappa \left[(\mathbf{Y}_e^\dagger \mathbf{Y}_e)^T \mathbf{C} + \mathbf{C} (\mathbf{Y}_e^\dagger \mathbf{Y}_e) \right] \quad (4.8)$$

where in the SM $\kappa = -3/2$ and $K = -3g_2^2 + 2\text{Tr} \left(3\mathbf{Y}_u^\dagger \mathbf{Y}_u + 3\mathbf{Y}_d^\dagger \mathbf{Y}_d + \mathbf{Y}_e^\dagger \mathbf{Y}_e \right) + 2\lambda$; the normalization of λ is fixed by the Higgs self interaction: $\mathcal{L}_{\text{self}} = -\frac{\lambda}{2}(H^\dagger H)^2$. In the MSSM, $\kappa = +1$ and $K = -6g_2^2 - 2g_Y^2 + 2\text{Tr} \left(3\mathbf{Y}_u^\dagger \mathbf{Y}_u \right)$. Since our initial conditions will be always given at the scale M , we have written eq. (4.8) top-down, i.e. with $16\pi^2 t = \ln(M/Q)$. The equation (4.8) is valid in any flavour basis, in particular in the basis in which the Yukawa matrix $\mathbf{Y}_e$ is diagonal.¹¹ In that basis, eq. (4.8) simplifies to

$$\frac{d}{dt}\mathbf{C}^{AB} = -K\mathbf{C}^{AB} - \kappa \left[y_{e_A}^2 \mathbf{C}^{AB} + \mathbf{C}^{AB} y_{e_B}^2 \right] \quad (4.9)$$

where $y_{e_A}^2$ are the eigenvalues of the hermitean matrix $\mathbf{Y}_e^\dagger \mathbf{Y}_e$. In this form it can be elegantly solved [34]:

$$\mathbf{C}(t) = I_K \mathcal{J} \mathbf{C}(0) \mathcal{J} \quad (4.10)$$

where $\mathcal{J} = \text{diag}(I_e, I_\mu, I_\tau)$ and

$$\begin{aligned} I_K &= \exp \left(- \int_0^t K(t') dt' \right), \\ I_{e_A} &= \exp \left(- \kappa \int_0^t y_{e_A}^2(t') dt' \right). \end{aligned} \quad (4.11)$$

Note that any zero in the initial matrix $\mathbf{C}^{AB}$ is preserved by the RG evolution and that the phases of the $\mathbf{C}^{AB}$ entries do not evolve [35]. We also note that, since $I_K I_e^2 \approx 1$, the experimental constraint (2.14), with $m_\nu^{ee} = (v^2/4M)\mathbf{C}^{11}(0)$ and $v^2 = v^2(M_Z)$, is renormalized negligibly.

Although the solution (4.10) to the RG equation for $\mathbf{C}^{AB}$ is simple, qualitative features of the running of the neutrino mass eigenvalues m_{ν_a} and of the MNS matrix are often masked by the diagonalization procedure. In that approach it has to be performed *after*

¹⁰The MSSM RGEs given in ref. [31] allow to treat also the case in which squarks and/or gluino are much heavier than sleptons, charginos and neutralinos so that the decoupling procedure [33] can be employed; there are then four different operators which mix with each other below the squark/gluino threshold. Above it one has (in the notation of ref. [31]) $c_1^{ab} = 2c_{12}^{ab} = 2c_{21}^{ab} = c_3^{ab} \equiv \mathbf{C}^{ab}$ and the four equations of ref. [31] merge into the one quoted here.

¹¹This is because the matrix $\mathbf{E}_L$ defined in eq. (3.4) does not evolve; the matrices $\varepsilon_{L,R}^E$ analogous to the ones defined in eq. (4.1) are identically zero because below the scale M , at which the operator (3.11) is generated, the RGE for the leptonic Yukawa coupling (B.1,B.5) do not depend on the couplings $\mathbf{Y}_\nu$, the only one that could change their matrix structure during the evolution.

the evolution. It is therefore useful to derive the RGE directly for the mass eigenvalues and mixing angles. This is done by using a trick similar as for the CKM matrix. One defines the auxiliary antihermitean matrix ε^ν by the equation

$$\frac{d}{dt}U = U\varepsilon^\nu \quad (4.12)$$

where U satisfies eq. (3.13). Differentiating the equality $C^a\delta^{ab} = (U^T\mathbf{C}U)^{ab}$ and requiring that its rhs remains a diagonal matrix gives us the matrix ε^ν . The only difference is that the resulting equation depends on $(\varepsilon^\nu)^T$ instead of $(\varepsilon^\nu)^\dagger$. Hence, one obtains two separate equations for $\text{Re}(\varepsilon^\nu)$ and $\text{Im}(\varepsilon^\nu)$:

$$\begin{aligned} \text{Re}(\varepsilon^\nu)^{ab} &= \kappa A_{ab} \text{Re} \left(\sum_A U^{Aa*} y_{e_A}^2 U^{Ab} \right) \\ \text{Im}(\varepsilon^\nu)^{ab} &= \kappa (A_{ab})^{-1} \text{Im} \left(\sum_A U^{Aa*} y_{e_A}^2 U^{Ab} \right) \end{aligned} \quad (4.13)$$

where

$$A_{ab} \equiv \frac{C^a + C^b}{C^a - C^b} = \frac{m_{\nu_a} + m_{\nu_b}}{m_{\nu_a} - m_{\nu_b}} \quad (4.14)$$

and of course $\text{Re}(\varepsilon^\nu)^{aa} = \text{Im}(\varepsilon^\nu)^{aa} = 0$. For conserved CP (U real), those equations have been first derived in ref. [36]. For a general complex matrix U they have been given in [37]. The formulae for dU/dt written there is valid in a general electroweak basis; it reduces to (4.13) after setting $P = \mathbf{Y}_e^\dagger \mathbf{Y}_e = P^\dagger$ and passing to the basis in which P is diagonal. Thus,

$$\frac{d}{dt}U^{Aa} = \kappa \sum_{b \neq a} U^{Ab} \left[A_{ba} \text{Re} \left(\sum_B U^{Bb*} y_{e_B}^2 U^{Ba} \right) + \frac{i}{A_{ba}} \text{Im} \left(\sum_B U^{Bb*} y_{e_B}^2 U^{Ba} \right) \right] \quad (4.15)$$

$$\frac{d}{dt}C^a = - \left(K + 2 \sum_A y_{e_A}^2 |U^{Aa}|^2 \right) C^a. \quad (4.16)$$

Eq. (4.15) gives directly the running of the angles θ_{12} , θ_{23} , θ_{13} [36] and the phases δ , α_a , φ_A parameterizing the matrix $U^{Aa} = e^{i\varphi_A} U_{\text{MNS}}^{Aa}$ [37]. In general, even if the phases φ_A are zero at some scale they will be generated during the evolution. However, it is easy to see that the differential equations for the parameters of the MNS matrix (2.10) do not depend on the φ_A . Indeed, from eq. (4.13) it follows that ε^ν does not depend on φ_A ; furthermore, since $\dot{U}^{Aa} = e^{i\varphi_A} [\dot{U}_{\text{MNS}}^{Aa} + i\dot{\varphi}_A U_{\text{MNS}}^{Aa}]$, all factors $e^{i\varphi_A}$ cancel out in eq. (4.12).

If two of the three eigenvalues, say m_{ν_a} and m_{ν_b} , are equal at some scale t there is the freedom in choosing the matrix $U(t)$, corresponding to the redefinition $U(t) \rightarrow \tilde{U}(t) =$

$U(t)R$ where R is a rotation in the ab plane. For the evolution, R has to be fixed by the condition

$$\text{Re}[\sum_A \tilde{U}^{Aa*} y_{e_A}^2 \tilde{U}^{Ab}](t) = 0 \quad (4.17)$$

so that ε^ν is nonsingular. This is particularly important for considering mass patterns with exact degeneracy at the scale M . The mixing matrix U is then ambiguous at that scale in the tree level approximation and becomes determined by quantum corrections, no matter how small. We shall illustrate this point in the next section. Note also that for conserved CP and neutrinos ν_a and ν_b of opposite CP parities such an ambiguity of rotations R does not exist. In such a case it is convenient to work with a real matrix U^{Aa} and with the neutrino masses of opposite signs (see the comments at the end of Section 3). The RG equation (4.15) is then nonsingular because A_{ab} , instead of being divergent, vanishes.

Eqs. (4.15) and (4.16) are very convenient for a qualitative discussion of the impact of the RG evolution directly on the neutrino masses and mixing angles. As we shall see, several physical effects are in this approach more transparent than in the approach based on evolving the matrix $\mathbf{C}^{AB}$.

4.3 Evolution of the neutrino masses

The effects of the RG quantum corrections on the neutrino mass eigenvalues are simple and, except for a few special cases, not very interesting. The solution to eq. (4.16) reads

$$m_{\nu_a}(t) = I_K \exp\left(-2\kappa \int_0^t y_{e_A}^2(t') |U^{Aa}(t')|^2 dt'\right) m_{\nu_a}(0). \quad (4.18)$$

Neglecting the small y_e and y_μ Yukawa couplings¹²

$$m_{\nu_a}^2(t) = m_{\nu_a}^2(0) I_K^2 \exp\left(-4\kappa \int_0^t y_\tau^2(t') |U_{3a}|^2(t') dt'\right) \quad (4.19)$$

Factors I_K and κ are different for the SM and the MSSM ($\kappa = -3/2$ and $+1$, respectively). We see that the possibility of some change in the mass pattern caused by the evolution resides solely in the differences in the mixing matrix elements U_{3a} and their RG running. In the MSSM where $y_\tau^2 \approx (\tan\beta/100)^2$, $t_Z \approx 0.12$ for $M = 10^{10}$ GeV and U_{3a}^2 typically varies between 0 and $1/4$ (except for U_{33}^2), the exponent is at most of order of $\epsilon \equiv y_\tau^2 \log(M/M_Z)/16\pi^2 \approx \tan^2\beta \times 10^{-5} < 2.5 \times 10^{-2}$ for $\tan\beta < 50$. In the SM, $y_\tau^2 \approx 10^{-4}$

¹²In the SM the Yukawa couplings are unambiguously determined at the electroweak scale by the charged lepton masses; in the MSSM $(y_{e_A}^2)_{\text{MSSM}} = (y_{e_A}^2)_{\text{SM}}/\cos\beta$ where $\tan\beta \equiv v_u/v_d$ can vary from a few up to ~ 50 .

and, consequently, $\epsilon \approx 10^{-5}$. We can then estimate the changes in the mass squared differences:

$$\Delta m_{ab}^2(t) \equiv m_{\nu_a}^2(t) - m_{\nu_b}^2(t) = \Delta m_{ab}^2(0) - (\eta_a m_{\nu_a}^2(0) - \eta_b m_{\nu_b}^2(0))\epsilon \quad (4.20)$$

where we have neglected I_K which is always close to 1, and the factors $\eta_{a(b)} > 0$ are typically in the range $0 - 2$, depending on the values of U_{3a} factors and their evolution. Taking (for definiteness) $\Delta m_{ab}^2(0) = 0$, we see that the evolution of $\Delta m_{ab}^2(t)$ is limited by $m_{\nu_a}^2(0)\epsilon$ or $m_{\nu_b}^2(0)\epsilon$, i.e. by the value of the larger mass.

We see that the RG evolution cannot change the pattern of masses. It may still be of importance for precision tests of various models, particularly those with partly degenerate or degenerate patterns. We recall here that such patterns have (at least approximately) degenerate two and three neutrinos, respectively, at the scale M . Matching the experimental Δm^2 and ΔM^2 needs then some fine-tuning between the initial values and the RG quantum corrections. That point can, however, be meaningfully discussed only for each concrete model.

The special cases are those with equal masses, $m_{\nu_1}^2 = m_{\nu_2}^2 \neq m_{\nu_3}^2$ or $m_{\nu_1}^2 = m_{\nu_2}^2 = m_{\nu_3}^2$ at the scale M . They are called in this review two-fold and three-fold degeneracies, respectively, to be distinguishable from the more general partly degenerate or degenerate patterns. The interesting question we shall discuss in Subsection 4.5 and Section 5 is whether only quantum corrections can then explain the observed mass squared differences.

In the next sections we shall discuss the RG quantum corrections to the neutrino mixing. We shall focus on interesting qualitative effects that depend only on the broad classification of the neutrino mass pattern, and do not depend on such details as whether the masses are equal or only approximately equal.

4.4 Mixing of two neutrinos

We first examine the mixing of two neutrinos, which was investigated in many papers [32, 38, 39, 40, 41]. Strictly speaking, it could be physically relevant only if the atmospheric neutrino anomaly was due to the ν_μ - ν_τ oscillations whereas the solar neutrino deficit resulted from the ν_e - ν_{sterile} oscillations (at present strongly disfavoured by the SNO and Superkamiokande data). Moreover the two 2×2 neutrino systems would have to be completely independent due to some particular texture of the U matrix. Nevertheless we will see, that the evolution of the θ_{23} angle in the 3×3 scenario is, in some cases very similar to the evolution of the mixing angle of two neutrinos only. It is therefore instructive to discuss the 2×2 evolution and to compare it later with the more realistic 3×3 mixing.

We begin with a real matrix $\mathbf{C}^{AB}$ and hence, a real U^{Aa} :

$$U^{Aa} = \begin{pmatrix} c_\vartheta & -s_\vartheta \\ s_\vartheta & c_\vartheta \end{pmatrix} \quad (4.21)$$

where $s_\vartheta \equiv \sin \vartheta$, $c_\vartheta \equiv \cos \vartheta$. We shall consider the same or different CP parities of the two neutrinos $m_{\nu_1} m_{\nu_2} > 0$ or $m_{\nu_1} m_{\nu_2} < 0$, respectively. From eq. (4.15) one finds the following equations for the mixing angle

$$\frac{ds_\vartheta}{dt} = \kappa A_{21} (y_{e_2}^2 - y_{e_1}^2) s_\vartheta c_\vartheta^2 \quad \frac{dc_\vartheta}{dt} = -\kappa A_{21} (y_{e_2}^2 - y_{e_1}^2) s_\vartheta^2 c_\vartheta, \quad (4.22)$$

where $\kappa = -3/2$ and $+1$ in the SM and the MSSM, respectively. Eqs. (4.22) give

$$\frac{d}{dt} \sin^2 2\vartheta = 2\kappa A_{21} (y_{e_2}^2 - y_{e_1}^2) \sin^2 2\vartheta \cos 2\vartheta \quad (4.23)$$

known from the literature¹³ [32, 38, 42, 43, 44]. Eq. (4.23) has a trivial fixed point (FP) at $\sin^2 2\vartheta = 0$ but, contrary to the statements made in some papers, the maximal mixing $\sin^2 2\vartheta = 1$ is not its FP: Although, naively, $\sin^2 2\vartheta(t) \equiv 1$ solves eq. (4.23), it is easy to see that $s_\vartheta(t) = \pm 1/\sqrt{2}$ does not solve eqs. (4.22).¹⁴

It is clear from eq. (4.22) that, for fixed values of the Yukawa couplings $y_{e_A}^2$ and for some fixed evolution “time” $t_Z \equiv (1/16\pi^2) \ln(M/M_Z)$, the evolution of the mixing angle depends on the factor $A_{21} = (m_{\nu_2} + m_{\nu_1})^2 / \Delta m_{21}^2$. It is always small when the two neutrinos have opposite CP-parities i.e. when $m_{\nu_1} m_{\nu_2} < 0$ (as follows from eq. (4.18), m_{ν_a} cannot change sign during the evolution) or if the neutrino mass spectrum is hierarchical ($m_{\nu_1}^2 \ll m_{\nu_2}^2$ or $m_{\nu_2}^2 \ll m_{\nu_1}^2$). The evolution of the mixing angle is then negligible [39, 35, 45, 34]. For $m_{\nu_1} m_{\nu_2} > 0$ the evolution of ϑ can be significant, particularly if the neutrino masses are nearly degenerate so that $|A_{21}\epsilon| \gg 1$ where $\epsilon \equiv (y_{e_2}^2 - y_{e_1}^2)t_Z$. It can be checked that for $|A_{21}|\epsilon \gtrsim 3$ the FP at $\sin^2 2\vartheta = 0$ is reached at the electroweak scale.

In view of the maximal ν_μ - ν_τ mixing needed to explain the atmospheric neutrino data, some attention was paid to the possibility of increasing ϑ by the RG corrections from a small value at the M scale to a (nearly) maximal ($|\vartheta| \approx \pi/4$) at the electroweak scale [32, 38, 39, 40, 41]. As we said earlier, the maximal mixing is not a FP of eqs. (4.22), so such an “explanation” of large mixing can merely be due to a coincidence of the running “time” and the initial values of the angles and the masses. The value of $|A_{21}|\epsilon$ must be

¹³Its rhs is more frequently written in the equivalent form $2\kappa \frac{C^{22} + C^{11}}{C^{22} - C^{11}} (y_{e_2}^2 - y_{e_1}^2) \sin^2 2\vartheta \cos^2 2\vartheta$.

¹⁴For example, for constant $\eta \equiv \kappa A_{21} (y_{e_2}^2 - y_{e_1}^2)$, eq. (4.22) is solved by $s_\vartheta^2(t) = s_0^2 / (s_0^2 + c_0^2 \xi)$ where $\xi \equiv \exp(-2\eta t)$ and s_0 is the initial value of s_ϑ . It is then straightforward to check that $\sin^2 2\vartheta(t) = \sin^2 2\vartheta_0 \xi / (s_0^2 + c_0^2 \xi)^2$ solves eq. (4.23) for any initial value $\sin^2 2\vartheta_0$. Thus, for $\sin^2 2\vartheta_0 = 1$, eq. (4.23) has two solutions: $\sin^2 2\vartheta(t) \equiv 1$ and the one given here but only the latter satisfies the underlying eq. (4.22).

in the range such that the evolution is non-negligible, but not strong enough to reach the FP. This can be most easily seen if we return to the solution (4.10) written in the form

$$\mathbf{C}^{AB}(t) = \begin{pmatrix} C^{11}(t) & C^{12}(t) \\ C^{12}(t) & C^{22}(t) \end{pmatrix} \propto \begin{pmatrix} C_0^{11} & C_0^{12} I_{21}(t) \\ C_0^{12} I_{21}(t) & C_0^{22} I_{21}^2(t) \end{pmatrix} \quad (4.24)$$

where $I_{21}(t) = I_{e2}/I_{e1} = \exp\left(-\kappa \int_0^t (y_{e2}^2 - y_{e1}^2)(t') dt'\right)$. One then has

$$\sin^2 2\vartheta(t) = \frac{4 [C^{12}(t)]^2}{[C^{11}(t) - C^{22}(t)]^2 + 4 [C^{12}(t)]^2} \quad (4.25)$$

It is obvious that $\sin^2 2\vartheta(t) \approx 1$ is obtained whenever $C^{11}(t) \approx C^{22}(t)$. Since $C^{22}(t)$ evolves differently from $C^{11}(t)$ (as $y_{e2}^2 \gg y_{e1}^2$) it is relatively easy to device the situation in which, at the initial scale M , $\sin^2 2\vartheta_0$ is small (this requires $|C_0^{11} - C_0^{22}| \gg 2|C_0^{12}|$) and the evolution is such that at some lower scale $C^{11}(t) = C^{22}(t)$ holds (this obviously requires $C_0^{22}C_0^{11} > 0$). With the judicious choice of the $\mathbf{C}^{AB}(0)$ matrix elements and $\epsilon \equiv (y_{e2}^2 - y_{e1}^2)t_Z$ it is possible to obtain $\sin^2 2\vartheta \approx 1$ at the electroweak scale [40, 41]. Expressing C_0^{11} and C_0^{22} in terms of the neutrino masses and mixing angle at the initial scale M , the relevant condition $C_0^{11} = C_0^{22} I_{21}^2(t_Z)$ reads [41]

$$(m_{\nu_1}^{(0)} c_{\vartheta_0}^2 + m_{\nu_2}^{(0)} s_{\vartheta_0}^2) = (m_{\nu_1}^{(0)} s_{\vartheta_0}^2 + m_{\nu_2}^{(0)} c_{\vartheta_0}^2) I_{21}^2(t_Z) \quad (4.26)$$

where $s_{\vartheta_0} \equiv \sin \vartheta(0)$ and $m_{\nu_a}^{(0)} \propto C^a(0)$. It is clear that for $|m_{\nu_2}^{(0)}| \approx |m_{\nu_1}^{(0)}|$ and $s_{\vartheta_0} \approx 0$ (or $c_{\vartheta_0} \approx 0$) satisfying the condition (4.26) requires $m_{\nu_1} m_{\nu_2} > 0$. On the other hand, for $m_{\nu_1} m_{\nu_2} < 0$ the product $C_0^{11} C_0^{22}$ can be positive only if $m_{\nu_2}^2 \ll m_{\nu_1}^2$ or $m_{\nu_2}^2 \gg m_{\nu_1}^2$, which leads to a hierarchy $C_0^{11} \gg C_0^{22}$ or $C_0^{11} \ll C_0^{22}$. Getting $C^{22}(t) = C^{11}(t)$ requires then $I_{21} \ll 1$ or $\gg 1$. Thus, for $m_{\nu_1} m_{\nu_2} < 0$ the evolution parameter ϵ must be large, too large to be accommodated in realistic theories. This is in agreement with our earlier observation that for opposite CP parities of the two neutrinos and/or their hierarchical masses the RG evolution is very weak.

Of course, since $\sin^2 2\vartheta = 1$ is not the FP of the RGE, $\sin^2 2\vartheta = 1$ can hold only at one particular scale.

Similar strategy can be also applied to analyze the general complex $\mathbf{C}^{AB}$. Once the parameters of the U^{Aa} matrix

$$U = \begin{pmatrix} e^{i\varphi_1} & 0 \\ 0 & e^{i\varphi_2} \end{pmatrix} \begin{pmatrix} c_{\vartheta} & -s_{\vartheta} \\ s_{\vartheta} & c_{\vartheta} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & e^{i\alpha_2} \end{pmatrix} \quad (4.27)$$

are expressed explicitly in terms of the $\mathbf{C}^{AB}$ entries, the solution (4.10) allows to obtain analytic formula for $\sin^2 2\theta(t)$ and to study its behaviour. The relevant formulae have been given in ref. [46] (see also [47]). As could be expected, for fixed initial values of $\sin^2 2\theta$ and

the neutrino masses $|m_{\nu_1}|$ and $|m_{\nu_2}|$, the $\sin^2 2\theta$ obtained by the RG evolution interpolates smoothly between its value obtained for $\alpha_2 = 0$ (i.e. $m_{\nu_1}m_{\nu_2} > 0$) and $\alpha_2 = \pm\pi/2$ (i.e. with $m_{\nu_1}m_{\nu_2} < 0$) [48].

4.5 Mixing of three neutrinos and fixed points

For conserved CP, it is straightforward to derive from (4.15) the equations for the three independent mixing parameters s_{12} , s_{23} and s_{13} [36, 37]. Neglecting y_e and y_μ Yukawa couplings we get, both in the SM and in the MSSM:

$$\begin{aligned}\dot{s}_{12} &= -c_{12}(s_{12}s_{23} - c_{12}c_{23}s_{13})(-c_{12}s_{23} - s_{12}c_{23}s_{13})\kappa A_{21}y_\tau^2 \\ &- s_{12}c_{12}c_{23}s_{13}(s_{12}s_{23} - c_{12}c_{23}s_{13})\kappa A_{31}y_\tau^2 \\ &+ c_{12}^2c_{23}s_{13}(-c_{12}s_{23} - s_{12}c_{23}s_{13})\kappa A_{32}y_\tau^2\end{aligned}\tag{4.28}$$

$$\begin{aligned}\dot{s}_{23} &= s_{12}c_{23}^2(s_{12}s_{23} - c_{12}c_{23}s_{13})\kappa A_{31}y_\tau^2 \\ &- c_{12}c_{23}^2(-c_{12}s_{23} - s_{12}c_{23}s_{13})\kappa A_{32}y_\tau^2\end{aligned}\tag{4.29}$$

$$\begin{aligned}\dot{s}_{13} &= -c_{12}c_{23}c_{13}^2(s_{12}s_{23} - c_{12}c_{23}s_{13})\kappa A_{31}y_\tau^2 \\ &- s_{12}c_{23}c_{13}^2(-c_{12}s_{23} - s_{12}c_{23}s_{13})\kappa A_{32}y_\tau^2.\end{aligned}\tag{4.30}$$

The evolution of the mixing angles can be classified into several universal types of behaviour, depending on the magnitude of the factors A_{ab} in eqs. (4.28-4.30). We note that, neglecting the small effects of mass evolution, all possible mass configurations in patterns *i*)-*iii*) listed at the end of Section 2 (except for the case of neutrinos (approximately) degenerate in mass and all having the same CP parity - to be discussed later) give one of the following four structures:

- a) $A_{31} \approx A_{32}$ and $|A_{31}| \approx |A_{21}| \approx 1$
- b) $A_{31} \approx A_{32}$ and $|A_{21}| \gg |A_{31}|, |A_{21}| \gg 1$
- c) $A_{32} \approx A_{21} \approx 0, |A_{31}| \gg 1$
- d) $A_{31} \approx A_{21} \approx 0, |A_{32}| \gg 1$

For hierarchical masses and for partly degenerate structure with opposite CP parities of the (almost) degenerate neutrinos all A_{ab} are $\mathcal{O}(1)$. For partly degenerate pattern with same CP parities or for degenerate pattern, at most one of them is large. In the first case, it follows from eqs. (4.28)-(4.30) and the value of ϵ ranging from 10^{-5} in the

SM to 2.5×10^{-2} in the MSSM with $\tan \beta \approx 50$ that the evolution of the angles is very weak [34, 54, 36, 37]. On the other hand, for one of the A_{ab} factors sufficiently big so that $|A_{ab}\epsilon| \gtrsim 1$, the angles evolve to an infrared quasi-fixed point (FP) [36, 37]. It is clear from eqs. (4.28)-(4.30) and from the parametrization (2.10) of the MNS matrix U , that depending on which A_{ab} is large, the fixed points are either at $U_{31} = 0$ or $U_{32} = 0$. Before discussing the approach to those fixed points in more detail, we can already now summarize several qualitative conclusions.

It is interesting to notice that in both fixed points we get the same relation between the mixing angles

$$\sin^2 2\theta_{12} = \frac{s_{13}^2 \sin^2 2\theta_{23}}{(s_{23}^2 c_{13}^2 + s_{13}^2)^2} \quad \text{with} \quad s_{23}^2 = \frac{1}{2} \left(1 \pm \sqrt{1 - \sin^2 2\theta_{23}} \right). \quad (4.31)$$

Thus, contrary to the 2×2 case, quantum corrections can give now an interesting, non-trivial fixed point relation. It is particularly interesting in the context of the present experimental indications for small θ_{13} angle from CHOOZ and maximal atmospheric and solar mixing ($\sin^2 2\theta_{23} \approx 1$ and $\sin^2 2\theta_{12} \approx 1$). The relation (4.31) is inconsistent with such a pattern of mixing. We stress that quantum corrections summarized in RGEs eqs. (4.28)-(4.30), if large, always give (4.31). Thus, if the presently most likely pattern of mixing is confirmed experimentally, all mass patterns generating large quantum corrections through RGE are ruled out!

Special cases easy to consider (still before a detailed study of the approach to the fixed points) are exact degeneracies at the scale M : $m_{\nu_1}^2 = m_{\nu_2}^2 \neq m_{\nu_3}^2$ or $m_{\nu_1}^2 = m_{\nu_2}^2 = m_{\nu_3}^2$. As follows from the discussion surrounding eq. (4.17), for the same CP parities, $m_{\nu_1} = m_{\nu_2}$ or $m_{\nu_1} = m_{\nu_3}$ or $m_{\nu_2} = m_{\nu_3}$ the angles must satisfy the FP relation (4.31) already at the scale M . If this is ruled out by experiment, then the two-fold degeneracy needs $m_{\nu_1} = -m_{\nu_2}$ and the three-fold degeneracy is ruled out. This conclusion holds (both in the SM and MSSM) under the assumption that the RGE corrections are the dominant ones (see next section for other possibilities). For $m_{\nu_1} = -m_{\nu_2}$ and $m_{\nu_1}^2 \gg m_{\nu_3}^2$, with $|m_{\nu_3}^2 - m_{\nu_2}^2| \approx \Delta M^2$, we can ask if quantum corrections can explain $\Delta m^2 = |m_{\nu_2}^2 - m_{\nu_1}^2|$. The answer to this question is positive in the MSSM [49] (in the SM see [50]) and will be discussed in more detail in Section 5. Another point worth a discussion is what if experiment will eventually be consistent with the FP relation (4.31). Two-fold degeneracy is easy: we need $m_{\nu_1} = m_{\nu_2}$, $|m_{\nu_3}^2 - m_{\nu_2}^2| \approx \Delta M^2$ and (as discussed in Section 5) quantum corrections can explain Δm^2 . With three-fold degeneracy, we necessarily have at least one pair of neutrinos with the same CP parities but the question is whether we can explain both Δm^2 and ΔM^2 by the discussed here class of quantum corrections. The answer is negative [51, 52, 53]. This can be explained as follows: from eq. (4.19) we would need

$|U_{31}| \approx |U_{32}|$ (to keep $m_{\nu_1}^2 \approx m_{\nu_2}^2$) during the entire evolution and, at the same time, $|U_{33}| - |U_{32}| \sim \mathcal{O}(1)$ to generate sufficiently large ΔM^2 . With $\theta_{13} \approx 0$ the first condition implies $\sin^2 2\theta_{12} \approx 1$ and the second $\sin^2 2\theta_{23} \approx 1$ ($|U_{33}| - |U_{32}|$ can be at most $\sim 1/2$). This is however incompatible with the relation (4.31) which should be satisfied! So, even if experiment was consistent with (4.31), the three-fold degenerate mass spectrum would still be unacceptable since ΔM^2 and Δm^2 could not be explained by quantum corrections (always under the assumption about the dominance of the RG corrections).

Let us now discuss the approach to the fixed point and concentrate first on the structure b) which can be realized for partly degenerate or degenerate patterns, with same CP parities of ν_1 and ν_2 . Since $A_{31} \approx A_{32}$ and $|A_{21}| \gg |A_{31}|$, equations (4.28-4.30) reduce to:

$$\begin{aligned}\dot{s}_{12} &= -c_{12}(s_{12}s_{23} - c_{12}c_{23}s_{13})(-c_{12}s_{23} - s_{12}c_{23}s_{13})A_{21}y_\tau^2 - c_{12}s_{23}c_{23}s_{13}A_{32}y_\tau^2, \\ \dot{s}_{23} &= s_{23}c_{23}^2A_{32}y_\tau^2, \\ \dot{s}_{13} &= c_{23}^2s_{13}c_{13}^2A_{32}y_\tau^2\end{aligned}\tag{4.32}$$

(we take MSSM, with $\kappa = 1$). The equation for $\dot{s}_{23}$ is the same as in the 2×2 scenario discussed in the previous subsection. The evolution of θ_{23} in the two cases is formally not identical as the evolution of the mass factor A_{32} depends now also on the remaining mixing angles θ_{12} and θ_{13} . Nevertheless, the qualitative behaviour of s_{23} is similar because in most cases the scale dependence of A_{32} can be neglected. Denoting

$$\xi_\tau \equiv \exp\left(-\int_0^\tau 2A_{32}(t')y_\tau^2(t')dt'\right) \approx \exp(-2A_{32}(0)\epsilon)\tag{4.33}$$

the solution for $s_{23}^2(t)$ reads

$$s_{23}^2(t) = s_{23}^2(0) / [s_{23}^2(0) + c_{23}^2(0)\xi_\tau]\tag{4.34}$$

and yields

$$\sin^2 2\theta_{23}(t) = \xi_\tau \sin^2 2\theta_{23}(0) / [s_{23}^2(0) + c_{23}^2(0)\xi_\tau]^2.\tag{4.35}$$

The solution for s_{13}^2 can also be given in a closed form:

$$s_{13}^2(t) = s_{13}^2(0) / \left\{ s_{13}^2(0) + c_{13}^2(0) [s_{23}^2(0) + c_{23}^2(0)\xi_\tau] \right\}.\tag{4.36}$$

Thus, since $|A_{32}| \approx 1$, the evolution of both, s_{23} and s_{13} is very weak [54, 36, 37]. For example, the effect of the running for $\sin^2 2\theta_{\text{atm}}$ is a 2.5% change for extreme value of $\tan \beta \approx 50$.

For $|A_{21}\epsilon| \gg 1$ (recall we consider $m_{\nu_{2(1)}}^2 > |\Delta m^2|$ and $m_{\nu_1}m_{\nu_2} > 0$), the evolution is towards one of the two approximate fixed points of the RG equation for s_{12} . One can

easily check, for instance, by considering the equation for $d \tan \theta_{12}/dt$, that for $A_{21} > 0$ (i.e. for $\Delta m^2 > 0$) the point $U_{31} = 0$ is the UV fixed point and $U_{32} = 0$ is the IR fixed point. For $A_{21} < 0$ (i.e. for $\Delta m^2 < 0$) the situation is reversed. It is also interesting to notice that in the limit $s_{13} = 0$ we can follow analytically the approach to the fixed points. In this approximation

$$\dot{s}_{12} = s_{12} c_{12}^2 s_{23}^2 A_{21} y_\tau^2 \quad (4.37)$$

and the solution for s_{12}^2 is of the form (4.34), with $s_{23}(c_{23}) \rightarrow s_{12}(c_{12})$ and¹⁵

$$\xi_\tau \rightarrow \xi'_\tau = \exp \left(- \int_0^t 2s_{23}^2(t') A_{21}(t') y_\tau^2(t') dt' \right). \quad (4.38)$$

For $A_{21} > 0$, in the top-down running, the factor $\xi' \rightarrow 0$ exponentially with decreasing the scale $Q \rightarrow M_Z$ i.e. for growing $t \propto A_{21} y_\tau^2 \log(M/Q)$ and, consequently, we obtain $s_{12}(t) = \pm 1$ (depending on its initial sign) and approach IR fixed point at $U_{32} = 0$. Changing the signs of the right hand sides of eqs. (4.32) one can see that in the bottom-up evolution we approach $s_{12}^2(t) \approx 0$ exponentially, i.e. the UV fixed point at $U_{31} = 0$. For $A_{21} < 0$ we get the reversed situation, in accord with our general expectations.

It is interesting to estimate the values of A_{21} and $\tan \beta$, for which the approach to the IR fixed points is seen. In ref. [36] we estimated that for approaching the fixed point during the evolution in the range (M, M_Z) with $M \approx 10^{10}$ GeV one needs $A_{21} \epsilon(M) > 3$, i.e. for $\tan \beta = 20$ one needs $m_{\nu_1} \approx m_{\nu_2} \gtrsim 10^{-4}$ eV for $\Delta m^2 \sim 10^{-10}$ eV² and $m_{\nu_1} \approx m_{\nu_2} \gtrsim 0.01$ eV for $\Delta m^2 \gtrsim 10^{-6}$ eV². The qualitative change from a negligible evolution to the FP behaviour at the electroweak scale is abrupt and occurs in the small range $0.5 \lesssim |A_{ab} \epsilon| \lesssim 3$.

Finally we note that, from the point of view of the initial conditions at the scale M , the UV fixed point looks not realistic as the neglected muon Yukawa coupling y_μ quickly destabilizes it during the evolution. We conclude that for hierarchical and partly degenerate mass patterns the evolution of the mixing angles is either very mild or shows (for $|A_{21}| \epsilon \gtrsim 3$) a fixed point behaviour.

The evolution of mixing angles in the degenerate case, $m_{\nu_3}^2 \approx m_{\nu_2}^2 \approx m_{\nu_1}^2 \sim \mathcal{O}(\Delta M^2)$ or larger, partly falls into the same classes of behaviour. Indeed, as long as $A_{31} \approx A_{32}$ with $|A_{31}| \lesssim \mathcal{O}(1)$ and $|A_{21}| \gg 1$, the angles evolve according to the same equations (4.32). One can easily identify the mass patterns of the degenerate case that fall into

¹⁵Eqs. (4.37) and (4.38), after solving for $s_{23}(t)$ and in the approximation of constant A_{21} and A_{23} , give:

$$s_{12}^2(t) = s_{12}^2(0) / \left\{ s_{12}^2(0) + c_{12}^2(0) [c_{23}^2(0) + s_{23}^2(0) \xi_\tau^{-1}]^{-A_{21}/A_{32}} \right\}.$$

this category: the necessary condition is that m_{ν_1} and m_{ν_2} are of the same sign. For the evolutions of s_{12} we then closely follow the two possibilities, depending on the sign of A_{21} , discussed for the partial degeneracy with $m_{\nu_1}m_{\nu_2} > 0$. We simply note that larger values of $|A_{21}|$ are generic for the present case and the approach to the fixed points is faster. However, the evolution of s_{23} and s_{13} are guaranteed to be mild only if m_{ν_1} and m_{ν_2} are negative. For positive m_{ν_1} and m_{ν_2} , i.e. when all masses have the same CP parity, we can have $|A_{32}\epsilon| \gtrsim 3$ for $\tan\beta > 40$ (due to the bound (2.14) $|A_{32}\epsilon| \gtrsim 3$ cannot be realized for $\tan\beta < 40$). The angle θ_{23} behaves then as the angle ϑ of the 2×2 mixing discussed in Section 4.3 and, according to the solution (4.34), s_{23} is exponentially focused to the stable FP $s_{23}(t) = 0$ or $s_{23}(t) = \pm 1$, depending on the sign of A_{32} , and on the direction of the evolution. The angle s_{13} behaves in a similar way except that, as follows from eq. (4.36), it does not reach the value $s_{13}^2 = 1$ when $s_{23}^2 \rightarrow 1$ (due to the presence of the factor c_{23}^2 in its RGE, the evolution of s_{13} is then “frozen”). We conclude that in the regime in which the approach to the fixed points is relevant, the pattern with approximately degenerate neutrino masses and the same all three CP parities is not acceptable.

As we said earlier, the approach to the FP behaviour is abrupt as a function of $|A_{31}\epsilon| \approx |A_{32}\epsilon|$. However, in the small transition region of the values of A_{31} and A_{32} , in agreement with our discussion in the previous subsection, it is possible to chose the initial condition for s_{23} so to get $\sin^2 2\theta_{23} = 1$, $s_{13} \approx 0$ at the electroweak scale. This was exploited in ref. [55] as a possible mean to obtain maximal atmospheric neutrino mixing from the initially small $\sin^2 2\theta_{23}$. Since $|A_{21}\epsilon|$ is always much larger than $|A_{32}\epsilon|$, the evolution of s_{12} is then such that the FP at $U_{32} = 0$ or $U_{31} = 0$ is quickly reached. Thus, a realistic solution, with maximal θ_{23} and $s_{13} \approx 0$, has $\sin^2 2\theta_{12} \approx 0$ and, moreover, the scheme cannot work unless there is an extreme fine tuning of the initial parameters [37].

The remaining degenerate mass patterns can be classified according to the relations $A_{32} \approx A_{21} \approx 0$ or $A_{31} \approx A_{21} \approx 0$. Consider first $A_{21} \approx A_{32} \approx 0$, i.e. $m_{\nu_1} \approx -m_{\nu_2} \approx m_{\nu_3}$. The equations for the evolution of the mixing angles can be approximated as

$$\begin{aligned}\dot{s}_{12} &= -s_{12}c_{12}c_{23}s_{13}(s_{12}s_{23} - c_{12}c_{23}s_{13})A_{31}y_\tau^2, \\ \dot{s}_{23} &= s_{12}c_2^2(s_{12}s_{23} - c_{12}c_{23}s_{13})A_{31}y_\tau^2, \\ \dot{s}_{13} &= -c_{12}c_{23}c_3^2(s_{12}s_{23} - c_{12}c_{23}s_{13})A_{31}y_\tau^2.\end{aligned}\tag{4.39}$$

These equations exhibit IR quasi-fixed point behaviour for $A_{31}\epsilon \ll -1$, corresponding to $U_{31} = 0$. As before, at the fixed point the angles satisfy the relation $s_{13} = \tan\theta_{12}\tan\theta_{23}$. Since $\dot{s}_{12}$ is proportional to s_{12} and suppressed by s_{13} , the running of s_{12} is weak. The IR fixed point is reached due to strong running of s_{23} and s_{13} . For $A_{31}\epsilon \gg 1$, $s_{23}(M_Z)$ is strongly focused at ± 1 . Thus, the mass and $\tan\beta$ configurations leading to $A_{31}\epsilon \gg 1$ are unacceptable. For $A_{31} \approx A_{21} \approx 0$ and $A_{32} < 0$ we get IR fixed point in $U_{32} = 0$.

The RG running of the mixing of three neutrinos for a complex matrix U , i.e. non-zero phase δ and $\alpha_1, \alpha_2 \neq 0$ modulo $\pi/2$ with all masses m_{ν_a} positive by definition, has been also investigated in the literature [37]. For initially degenerate ν_a and ν_b the mixing pattern is determined by the condition $\text{Re}(U^{3a}U^{3b*}) \equiv \text{Re}(U_{\text{MNS}}^{3a}U_{\text{MNS}}^{3b*}) = 0$ playing in the general complex case the role of the FP condition. Qualitatively, our conclusions remain unchanged also in this case. For example, starting with $m_{\nu_1}^2 = m_{\nu_2}^2$, $\sin^2 2\theta_{23} = 1$, $s_{13} \approx 0$ and arbitrary phases α_1, α_2 at the scale M , the RG evolution leads to the relation [37]

$$\sin^2 2\theta_{12}(M_Z) = \sin^2 2\theta_{12}(M) \sin^2(\alpha_1 - \alpha_2) + \mathcal{O}(s_{13}^2) \quad (4.40)$$

We see therefore that maximal solar mixing can again be obtained only if the change of θ_{12} during the running is negligible, i.e. for $\alpha_1 - \alpha_2 \approx \pi/2$ at the scale M .

The running of the masses is always given by eq. (4.18) and is similar as for a real matrix U . The conclusion that the triple degeneracy of neutrinos at the scale M , $|m_{\nu_1}| = |m_{\nu_2}| = |m_{\nu_3}|$, cannot lead to an acceptable pattern of masses and mixing also remains valid.

In summary, with all $|A_{ab}\epsilon| \lesssim 0.5$ the evolution of the mixing is negligible. For the mass configurations such that at least one $|A_{ab}| \gg 1$ and $|A_{ab}\epsilon| \gtrsim 3$ the infrared fixed points are reached during the evolution, independently of further details of the mass matrices. However, only for $|A_{21}\epsilon| \gtrsim 3$ and $|A_{31}|, |A_{32}| \lesssim 1$, or for $A_{31}(A_{32})\epsilon \ll -3$ and $A_{31}(A_{32}) \approx A_{21} \approx 0$ the evolution is consistent with a large atmospheric mixing angle at low energy. The mass configurations $m_{\nu_1} \approx m_{\nu_2} \approx m_{\nu_3}$ and $\pm m_{\nu_1} \approx \mp m_{\nu_2} \approx m_{\nu_3}$ with $A_{31}(A_{32}) > 0$ also lead to the infrared fixed points but at the same time the atmospheric mixing angle converges to zero. We also note that for $|A_{21}\epsilon| \gtrsim 3$ only s_{12} runs to assure the fixed point relation, so the initial values for s_{23} and s_{13} have to be close to their experimental values already at the scale M . For $A_{31}\epsilon \lesssim -3$ or $A_{32}\epsilon \lesssim -3$, s_{23} and s_{13} evolve strongly and the evolution of s_{12} is weak. The IR fixed point relation (4.31) is always one equation for three angles. Insisting on a large atmospheric mixing angle, it correlates small (as follows from CHOOZ) θ_{13} angle with a small solar mixing angle. The fixed point solution makes the low energy angles dependent on only two, instead of in general three, initial conditions at the scale M . We conclude that quantum corrections encoded in the RG running of mixing angles may have dramatic impact on their physical values if the mass pattern is partial degeneracy or degeneracy. If the bimaximal mixing solution was confirmed, all the mass patterns leading to the FP would be ruled out, unless the low energy threshold corrections change the results.

5 Low energy threshold corrections

5.1 Threshold corrections in the SM

The Wilson coefficient $\mathbf{C}^{AB}$ of the dimension 5 $\Delta L = 2$ operator (3.11) is a renormalized parameter of the effective theory Lagrangian. Integrating its RGEs from the high scale M down to some scale $Q \approx M_Z$ allows to resum potentially large corrections involving $\ln(M/Q)$ to all orders of the perturbation expansion. However, since the low energy scale Q is not a priori determined by any physical requirement (apart from the condition $Q \approx M_Z$), the neutrino masses and mixing angles computed in the tree-level approximation from the Wilson coefficient $\mathbf{C}^{AB}(Q)$ do depend (albeit weakly) on the actual choice of Q . This dependence can be removed by computing masses and mixing angles in the one-loop approximation in the $\overline{\text{MS}}$ scheme and with Q as the renormalization scale.

In general, to compute the physical neutrino masses and mixing one has first to perform the RG evolution of the entire matrix U and neutrino masses from the scale M down to the scale M_Z , include threshold corrections and subsequently re-diagonalize the resulting mass matrix $m_{ab}^{1\text{-loop}}$ by an additional unitary matrix U' . The physical matrix U_{MNS} (whose elements are probed in neutrino experiments) is then given as $e^{i\varphi_A} U_{\text{MNS}}^{Aa} = (U \cdot U')^{Aa}$ where U is the matrix obtained from the RG evolution.

In the basis in which neutrino masses are diagonal, we have

$$m_{ab}^{1\text{-loop}} = m_{\nu_a} \delta^{ab} + m_{\nu_a} I^{ab} + m_{\nu_b} I^{ba} \quad (5.1)$$

where

$$I^{ab} = \sum_{A,B} U^{Aa*} I_{AB}^{\text{th}} U^{Bb} \quad (5.2)$$

In the SM one finds [56]

$$I_{AB}^{\text{th}} = \frac{\delta^{AB}}{16\pi^2} \frac{g_2^2}{2} \frac{m_{e_A}^2}{M_W^2} \left[\frac{11}{8} - \frac{3}{2} \ln \frac{M_W}{Q} + \mathcal{O}(x_A \ln x_A) \right] + \dots \quad (5.3)$$

where $x_A \equiv m_{e_A}^2/M_W^2$. Note that since $g_2^2 m_{e_A}^2/2M_W^2 = y_{e_A}^2$, the coefficient of $\ln(1/Q)$ agrees with the coefficient κ of the $y_{e_A}^2 \mathbf{C}^{AB} + \mathbf{C}^{AB} y_{e_B}^2$ term in the RGE (4.8) for the SM. This confirms the correctness of the recent re-derivation [29] of the SM RGE. The remaining corrections indicated by dots in (5.3) affect only the overall scale of the neutrino masses and therefore are not interesting in view of the unspecified magnitude of the mass M in eq. (3.13).

As could be expected, in the SM the nontrivial part (5.3) of the low energy threshold corrections that changes the structure of the mass matrix, and hence of the matrix U^{Aa} , is

small and always negligible compared to the quantum effects described by the RG running, which are enhanced by large logarithm of the ratio M/M_Z . The correction (5.3) can be most easily taken into account by stopping the RG evolution of the Wilson coefficient C^{AB} at the scale $Q = M_W e^{-11/12}$.

5.2 Threshold corrections in the MSSM

In the supersymmetric model, apart from stabilizing the results obtained from the RG analysis, the low energy threshold corrections can be as or more important than the RG evolution and may have very important physical consequences for the neutrino masses and mixing angles [61, 60, 58]. Before we discuss some physical examples, it is worthwhile to adapt our calculational procedure to the new situation, so that the RG evolution effects and the threshold corrections can be treated on equal footing. Although the general procedure outlined earlier is in principle correct irrespectively of the relative magnitude of the RG and threshold effects, in practice it can mask simple qualitative features if the re-diagonalization due to threshold corrections is not a small perturbation.

We observe that, to a very good accuracy, in the solution (4.10) to the RGE the factors I_{e_A} (4.11) can be approximated by

$$I_{e_A} \approx 1 - \kappa \int_0^{t_Z} y_{e_A}^2 dt' \equiv 1 - I_A^{\text{rg}}. \quad (5.4)$$

With this approximation (which can fail only for unrealistically large values of $\tan \beta$) all quantum effects of the physics below the scale M , the RG running as well as the low energy threshold corrections, can be described by a single formula (5.1). The m_{ν_a} and the matrix U are now the neutrino mass eigenvalues and the neutrino mixing matrix, respectively, at the scale M , and the factors I^{ab} are given by eq. (5.2) with

$$I_{AB}^{\text{th}} \rightarrow I_{AB}^{\text{th}} - \delta^{AB} I_A^{\text{rg}} \equiv I_{AB}. \quad (5.5)$$

Again, the case of a real matrix U is particularly easy because then the formula (5.1) can be written as

$$m_{ab}^{1\text{-loop}} = m_{\nu_a} \delta^{ab} + (m_{\nu_a} + m_{\nu_b}) I^{ab} \equiv m_{\nu_a} \delta^{ab} + \Delta m_{ab} \quad (5.6)$$

with the right hand side symmetric and real i.e. Hermitean. One can then use the formal perturbation calculus of quantum mechanics (see e.g. [57]) to find corrections to neutrino masses and mixing angles. Of course, if the threshold corrections are absent¹⁶ one has $I_{AB} \approx \delta^{AB} I_A$ with $|I_\tau| \gg |I_\mu| \gg |I_e|$.

¹⁶If the threshold corrections are universal, i.e. if $I_{AB}^{\text{th}} = I^{\text{th}} \delta^{AB}$, they can be absorbed into the overall scale of neutrino masses and do not influence neither the mixing angles nor the ratio $\Delta M^2 / \Delta m^2$.

Several physically interesting situations can be discussed. In the following we shall mainly focus on the scenarios with equal three or at least two masses at the scale M . Although the discussion of the mixing angles does not depend on whether the masses are equal or only approximately equal, the former case is much more constrained for the masses themselves and therefore more interesting.

5.2.1 Three-fold degeneracy and flavour diagonal corrections

In the MSSM the complete formulae for I_{AB}^{th} are lengthy due to additional contributions of $H^\pm$ and chargino/charged slepton and neutralino/sneutrino loops [56] but we do not need them for our discussion. First of all, it is interesting to investigate whether it is possible to generate the right mass squared differences starting from degenerate neutrinos at the scale M : $|m_{\nu_1}| = |m_{\nu_2}| = |m_{\nu_3}|$ (we again allow for different CP parities of the neutrinos). From the arguments presented in Section 4 it follows that this is impossible if the factors I_{AB} are dominated by the RG effects, i.e. when $I_{AB} \approx \delta^{AB} I_A$ with $|I_\tau| \gg |I_\mu|, |I_e|$. The degeneracy Ansatz is however potentially interesting as it might help to understand the bimaximal mixing. Also, the neutrino masses of order of few eV are necessary if the neutrinos are to play a role of the hot component of the Dark Matter.¹⁷ It is therefore interesting to see if some other corrections can split the mass squares.

We start with the flavour diagonal threshold corrections: $I_{AB} = \delta^{AB} I_A$ and consider first the mass pattern $m_{\nu_1} = -m_{\nu_2} = m_{\nu_3} \equiv m_\nu$. The formula (5.6) now takes the form

$$m_{ab}^{1\text{-loop}} = m_\nu \begin{pmatrix} 1 + 2U_{A1}^2 I_A & 0 & 2U_{A1} U_{A3} I_A \\ 0 & -1 - 2U_{A2}^2 I_A & 0 \\ 2U_{A1} U_{A3} I_A & 0 & 1 + 2U_{A3}^2 I_A \end{pmatrix} \quad (5.7)$$

The corrections to neutrino masses can be calculated perturbatively. Because $m_{\nu_1} = m_{\nu_3}$, one applies here the perturbation calculus to the case with degeneracy of the unperturbed “energy levels” [57]. Instead of solving the “secular” equation for the corrected eigenvalues, it is however better to exploit the freedom of an arbitrary rotation in the plane (13): $U \rightarrow UR_{13}$. This freedom can be used to diagonalize the perturbation by requiring

$$\sum_A U_{A1} U_{A3} I_A = 0 \quad (5.8)$$

which fixes the matrix U after taking into account the perturbation. The correction Δm_{ab} to the zeroth order neutrino mass matrix becomes then diagonal and the neutrino masses

¹⁷The hot component of the Dark Matter was previously needed to account for formation of largest scale structures in the Universe. At present, in view of the observational evidence for a significant contributions of the cosmological constant (or another form of dark energy as e.g. quintessence) to the energy density of the Universe, the hot component of the Dark Matter seems no longer necessary (but is not excluded).

are given by

$$|m_{\nu_a}| = m_\nu \left(1 + 2U_{Aa}^2 I_A\right). \quad (5.9)$$

Let us now assume [58] that $I_e \neq 0$, $I_\mu = I_\tau = 0$. The condition (5.8) reduces then to $U_{11}U_{13} = 0$. The experimentally viable solution is $U_{13} = 0$, i.e. $s_{13} = 0$. Note that this relation replaces the FP relation (4.31) of Section 4. For the neutrino masses one finds

$$\Delta m_{21}^2 = -4m_\nu^2 \cos^2 2\theta_{12} I_e, \quad \Delta m_{32}^2 = -4m_\nu^2 s_{12}^2 I_e. \quad (5.10)$$

It is therefore possible to generate $\Delta M^2 \gg \Delta m^2$ provided the solar mixing is nearly maximal. The same result is obtained starting from $-m_{\nu_1} = m_{\nu_2} = m_{\nu_3} \equiv m_\nu$. Thus, if the underlying theory gives one of these two degeneracies together with the bimaximal mixing pattern, the low energy threshold corrections can be responsible for assuring $s_{13} = 0$ and generating the correct mass squared differences.

Of course it is unrealistic to expect only one correction I_A to be nonzero. Note however, that the results for the mixing angles and the ratio $\Delta M^2/\Delta m^2$ are not altered by shifting all corrections I_A by an overall additive constant: $I_A \rightarrow \tilde{I}_A \equiv I_A - I$. Indeed, using unitarity of the matrix U , the off-diagonal entries in the rhs of eq. (5.6) can be always written as $\sum_A U_{Aa}U_{Ab}\tilde{I}_A$ ($a \neq b$) with arbitrary I . For the diagonal entries instead one has $1 + 2\sum_A U_{Aa}^2 I_A = 1 + 2I + 2\sum_A U_{Aa}^2 \tilde{I}_A \approx (1 + 2I)(1 + 2\sum_A U_{Aa}^2 \tilde{I}_A)$. Hence, up to higher order terms, eq. (5.6) can be rewritten with $\tilde{I}^{ab} = \sum_A U_{Aa}U_{Ab}\tilde{I}_A$ and m_ν replaced by $\tilde{m}_\nu \equiv (1 + 2I)m_\nu$. Thus, $I_\mu = I_\tau \neq 0$ is equivalent to $I_\mu = I_\tau = 0$.

It follows that $|I_e| \gg |I_\tau|$, $|I_\mu|$, equivalent to $|\tilde{I}_e| \gg |\tilde{I}_\tau| \neq 0$, $\tilde{I}_\mu = 0$, should give only small correction to the result (5.10) and $s_{13} = 0$. For $m_{\nu_1} = -m_{\nu_2} = m_{\nu_3}$, solving eq. (5.8) (with $I_e = \tilde{I}_e$, $I_\mu = 0$ and $I_\tau = \tilde{I}_\tau$) one finds [58] that

$$\begin{aligned} s_{13} &= -\frac{s_{12}}{c_{12}} s_{23} c_{23} r + \mathcal{O}(r^2) \\ \Delta m_{32}^2 &\approx -4\tilde{m}_\nu^2 \tilde{I}_e s_{12}^2 \\ \Delta m_{21}^2 &\approx -4\tilde{m}_\nu^2 \tilde{I}_e \left[\cos 2\theta_{12} (1 - s_{23}^2 r) + (1 + 2c_{12}^2) s_{13}^2 \right] \end{aligned} \quad (5.11)$$

where $r \equiv \tilde{I}_\tau/\tilde{I}_e$. Thus, obtaining small but non-zero angle θ_{13} is also possible. To obtain the experimentally favoured value of ΔM^2 one must have $|\tilde{m}_\nu^2 \tilde{I}_e| \approx 1.6 \times 10^{-3} \text{ eV}^2$ i.e. $|I_e - I_\mu| \sim 10^{-3}$ and $I_\mu \approx I_\tau$ for m_ν in the eV range. In the MSSM the required hierarchy $|I_e| \gg |I_\tau|$, $I_\tau \approx I_\mu$ can arise from the low-energy threshold corrections only as a result of non-universality of the left-handed charged slepton masses (and sneutrino masses).¹⁸

¹⁸In the $\tilde{W}^\pm$ loop approximation to I_{AB}^{th} it was estimated in ref. [58] that in order to get $\tilde{I}_e \sim 10^{-3}$ and positive one needs $M_{\tilde{e}_L} \approx 1.7M_{\tilde{\mu}_L}$ (and $M_{\tilde{\mu}_L} \approx M_{\tilde{\tau}_L}$). However, from the full expression for I_{AB}^{th} it follows [56] that $\tilde{I}_e \sim 10^{-3}$ and positive can be achieved only for relatively light first chargino ($m_{C_1} \lesssim 300 \text{ GeV}$) and $M_{\tilde{\mu}_L} \approx (1.2 - 1.6)M_{\tilde{e}_L}$. For $M_{\tilde{e}_L} \approx (1.2 - 1.6)M_{\tilde{\mu}_L}$, which can be realized in the inverted hierarchy models [59], one gets $\tilde{I}_e \sim -10^{-3}$ i.e. positive Δm_{21}^2 for (almost) maximal solar mixing.

Moreover, for exactly maximal solar mixing, obtaining Δm^2 appropriate for the LAMSW solution requires $s_{13}^2 \approx 10^{-(2-3)}$. This is consistent with the CHOOZ data and is realized for $r \lesssim 0.1$, i.e. $|I_\tau - I_\mu| \lesssim 10^{-4}$. For $\tan \beta \lesssim 3$ such a small difference between the corrections I_τ and I_μ can be due to the RG effects in I_τ^{rg} and does not require any mass splitting between μ_L and τ_L . For larger values of $\tan \beta$, for which $I_\tau^{\text{rg}} - I_\mu^{\text{rg}} > 10^{-4}$, some conspiracy between $I_\tau^{\text{th}} - I_\mu^{\text{th}}$ due to $M_{\tilde{\mu}_L} < M_{\tilde{\tau}_L}$ and $I_\tau^{\text{rg}} - I_\mu^{\text{rg}}$ is required. The amount of the necessary fine tuning grows, of course, with $\tan \beta$. For the VO solution to the solar neutrino problem, with $\sin^2 2\theta_{12} = 1$, a more severe fine tuning is always necessary: $\Delta m^2 \sim \mathcal{O}(10^{-10} \text{ eV}^2)$ requires $|I_\tau - I_\mu| \lesssim 10^{-7}$ which is always smaller than $I_\tau^{\text{rg}} - I_\mu^{\text{rg}} \gtrsim 10^{-5}$. Thus obtaining the correct Δm^2 for this solution requires a cancellation at least to one part per hundred between the RG and threshold effects. For $\Delta m^2 \sim \mathcal{O}(10^{-10} \text{ eV}^2)$ one gets $s_{13}^2 \lesssim 10^{-8}$. Similar results are obtained also for $-m_{\nu_1} = m_{\nu_2} = m_{\nu_3}$.

It can be checked [60] that, for the initially degenerate neutrino masses, the patterns $m_{\nu_1} = -m_{\nu_2} = m_{\nu_3}$ or $-m_{\nu_1} = m_{\nu_2} = m_{\nu_3}$ and $|I_e| \gg |I_\tau|$, $I_\tau \approx I_\mu$ are the only ones that can produce the required mass squared splittings with flavour diagonal threshold corrections. The necessary condition is that the underlying theory valid above the scale M gives the bimaximal mixing in the basis in which eq. (5.8) is satisfied [58, 60].

5.2.2 Three-fold degeneracy and flavour non-diagonal corrections

Qualitatively new possibilities for mixing angles and for splitting initially degenerate neutrinos originate from nonzero off-diagonal elements of the corrections I_{AB}^{th} in eq. (5.2) [61, 60]. This is possible in the MSSM if the slepton mass matrices are not diagonal in the flavour space in the basis in which the leptonic Yukawa couplings are diagonal. The amount of flavour mixing is then best quantified in terms of the so-called mass insertions [62] defined as the ratio of the off-diagonal (in flavour space) elements of the charged sleptons mass squared matrices to some average charged slepton mass squared.¹⁹ Current limits on such mass insertions following from the non-observation of the decays $\mu \rightarrow e\gamma$ etc. are not very stringent: Only the insertion δ_{LR}^{12} mixing left(right)-handed $\tilde{\mu}$ with right(left)-handed $\tilde{e}$ has to be smaller than $\mathcal{O}(10^{-5})$. The insertions δ_{LR}^{13} and δ_{LR}^{23} causing similar, chirality changing, $\tilde{e} \leftrightarrow \tilde{\tau}$ and $\tilde{\mu} \leftrightarrow \tilde{\tau}$ transitions are bounded by ≈ 0.5 and ≈ 0.1 , respectively, for slepton masses $\sim 500 \text{ GeV}$. Bounds on the chirality preserving insertions δ_{LL}^{AB} , δ_{RR}^{AB} exist only for the $\tilde{e} \leftrightarrow \tilde{\mu}$ transition and are ≈ 0.2 for $M_{\tilde{l}} \sim 500 \text{ GeV}$. The other chirality preserving mass insertions are practically unrestricted.

¹⁹Due to the underlying $SU_L(2)$ symmetry, mass insertions inducing transitions between left-handed sleptons of different generations are always accompanied by the insertions inducing similar transitions between sneutrinos.

If the mass insertions are non-vanishing, there is a flavour non-diagonal contribution to I_{AB}^{th} . In general it takes the form

$$I_{AB}^{\text{th}} \approx \frac{1}{16\pi^2} \sum_{X,Y=L,R} \delta_{XY}^{AB} h_{XY}(M_{\tilde{l}_{\text{av}}}, m_{C_j}, m_{N_i}) \quad (5.12)$$

where h_{LL} , h_{RR} and h_{LR} are some functions of the chargino and neutralino masses m_{C_j} , m_{N_i} and of some average mass $M_{\tilde{l}_{\text{av}}}$ of charged sleptons. The largest values of I_{AB}^{th} can be obtained for relatively light charginos, heavy sleptons (~ 1 TeV) and $M_2/\mu \approx -1$ and reach $h_{LL}/16\pi^2 \lesssim \text{few} \times 10^{-3}$ (h_{RR} and h_{LR} are always smaller) [56]. For comparable chargino and slepton masses one has $h_{LL}/16\pi^2 \lesssim 2 \times 10^{-4}$. In principle the mass insertion approximation should fail for $|\delta_{XY}^{AB}| \lesssim 0.1$. In practice it works as an order of magnitude estimate even for $|\delta_{XY}^{AB}| \lesssim 1$ (the error is then of order 25%).

In the presence of non-zero mass insertions and for $m_{\nu_a} = m_{\nu_b}$ ($= -m_{\nu_c}$) the condition for vanishing of the appropriate off-diagonal entry of the correction to the zeroth order mass matrix reads

$$\sum_{A,B} U_{Aa} U_{Bb} I_{AB} = 0. \quad (5.13)$$

Consider first the situation in which the single correction $I_{e\mu}^{\text{th}}$, $I_{e\tau}^{\text{th}}$ or $I_{\mu\tau}^{\text{th}}$ dominates over all other corrections. The condition (5.13) gives then relations between the mixing angles that are different from (4.31) and are listed in Table 1 [61]. Only three of the nine possibilities are compatible with the bimaximal mixing: dominant $I_{\mu\tau}^{\text{th}}$ for $m_{\nu_1} = m_{\nu_3}$ or $m_{\nu_2} = m_{\nu_3}$ and dominant $I_{e\tau}^{\text{th}}$ for $m_{\nu_1} = m_{\nu_2}$. For initially degenerate neutrinos, the latter combination gives wrong relation $\Delta m^2 \approx 2\Delta M^2$. (Other six combinations leading through (5.13) to $\sin^2 2\theta_{12} \sim \sin^2 2\theta_{13}$ also give bad relations $\Delta m^2 \approx \Delta M^2$ or $\Delta M^2 = 0$.) The former two are however interesting giving $\Delta m^2 \approx -4m_\nu^2 \cos 2\theta_{12} \sin 2\theta_{23} I_{\mu\tau}^{\text{th}}$ and $\Delta M^2 \approx 4m_\nu^2 (1 + c_{12}^2) \sin 2\theta_{23} I_{\mu\tau}^{\text{th}}$. Obtaining $\Delta M^2 \approx 3.2 \times 10^{-3} \text{ eV}^2$ is therefore possible but only for $m_\nu \gtrsim 1 \text{ eV}$ and $\delta_{LL}^{23} \gtrsim 0.5$ i.e. for rather large flavour mixing in the slepton mass matrices.

Similarly as for a non-zero $\tilde{I}_e$ correction, the solar mass squared difference $\Delta m^2 \ll \Delta M^2$ can be generated either by an appropriately tuned departure of the angle $|\theta_{12}|$ from maximal value $\pi/4$ or by another, hierarchically smaller, correction $I \neq 0$. It has been demonstrated [60], that including on the top of the dominant $I_{\mu\tau}^{\text{th}}$ correction a hierarchically smaller perturbation in the form of either $\tilde{I}_\mu$, $\tilde{I}_\tau$ ($\tilde{I}_e$ does not work) or $I_{e\mu}^{\text{th}}$, $I_{e\tau}^{\text{th}}$ allows to split $m_{\nu_1} = -m_{\nu_2}$ even for exactly bimaximal mixing.²⁰

²⁰We note [60], that with a large I_e diagonal perturbation discussed in the Subsection 5.2.1, $\Delta m^2 \ll \Delta M^2$ (and $0 \neq s_{13}^2 \ll 1$) can be also induced by $I_{e\mu}^{\text{th}}$ or $I_{e\tau}^{\text{th}}$ instead of $I = \tilde{I}_\tau$ (small $I_{\mu\tau}^{\text{th}} \neq 0$ would

Table 1: Relations of the FP type between the mixing angles for dominant correction I_{AB}^{th}

	$m_{\nu_1} = m_{\nu_2}$	$m_{\nu_1} = m_{\nu_3}$	$m_{\nu_2} = m_{\nu_3}$
$I_{e\mu}^{\text{th}}$	$s_{13} = \cot 2\theta_{12} \cot \theta_{23}$	$\frac{s_{13}^2}{1-2s_{13}^2} = \cot \theta_{12} \tan \theta_{23}$	$\frac{s_{13}^2}{1-2s_{13}^2} = -\tan \theta_{12} \tan \theta_{23}$
$I_{e\tau}^{\text{th}}$	$s_{13} = -\cot 2\theta_{12} \tan \theta_{23}$	$\frac{s_{13}}{1-2s_{13}^2} = -\cot \theta_{12} \cot \theta_{23}$	$\frac{s_{13}}{1-2s_{13}^2} = \tan \theta_{12} \cot \theta_{23}$
$I_{\mu\tau}^{\text{th}}$	$\frac{2s_{13}}{1+s_{13}^2} = \tan 2\theta_{12} \tan 2\theta_{23}$	$s_{13} = -\tan \theta_{12} \cot 2\theta_{23}$	$s_{13} = \cot \theta_{12} \cot 2\theta_{23}$

In all those cases of two hierarchically different corrections the important difference between the two alternatives: perturbation by a diagonal correction $\tilde{I}_A$ or perturbation by an off-diagonal correction I_{AB}^{th} is that, for exactly bimaximal mixing, in the former case $\Delta m^2 \propto r^2$, whereas in the latter $\Delta m^2 \propto r$ only ($s_{13} \sim r$ in all cases), where $r \ll 1$ is the ratio of the smaller to larger correction as in (5.11) [60]. Therefore, similarly as for $|\tilde{I}_e| \gg |\tilde{I}_\tau|$, obtaining Δm^2 appropriate for the VO solution with $I_{\mu\tau}^{\text{th}}$ dominance and hierarchically smaller $I_{e\tau}$ or $I_{e\mu}$ would require some tuning of slepton masses to cancel too large a contribution of I_τ^{g} to I_τ .

Finally, we remind the reader that the relations listed in Table 1 remain approximately valid when the equalities of the masses are relaxed and replaced by the corresponding approximate degeneracies. The listed relations play then the role of the FP relations discussed in Section 4. They are satisfied at the electroweak scale irrespectively of the initial values of the angles. The observed Δm^2 and ΔM^2 can be obtained by adjusting the initial values of only approximately equal masses. The role of the threshold corrections I_{AB}^{th} is then the same as the role of the RG corrections. If large enough, they give one of the “fixed points” relations of Table 1.

5.2.3 Two-fold degeneracy and threshold corrections

We can also discuss the effect of threshold corrections in the case of the two-fold degeneracy $m_{\nu_1}^2 = m_{\nu_2}^2 \equiv m_\nu^2 \gg m_{\nu_3}^2$ or $m_{\nu_1}^2 = m_{\nu_2}^2 \ll m_{\nu_3}^2$. For $m_{\nu_1} = -m_{\nu_2}$ the Δm_{12} off-diagonal entry in eq. (5.6) automatically vanishes but the correction matrix Δm_{ab} as the whole needs not be diagonal. In the first order of the perturbation calculus [57], the neutrino masses are then given by eq. (5.9) and receive also further corrections of order $\mathcal{O}((\Delta m_{ab})^2/\max(|m_{\nu_3}|, |m_\nu|))$. The mixing angles also receive corrections of order $\mathcal{O}((\Delta m_{ab})^2/\max(|m_{\nu_3}|, |m_\nu|))$. This leads to $s_{13} \approx -rs_{23}(c_{23})$ ($r \equiv I_{e\mu(\tau)}^{\text{th}}/\tilde{I}_e$) and $\Delta m^2 \approx 4m_\nu^2 \sin 2\theta_{12} c_{23}(s_{23}) I_{e\mu(\tau)}^{\text{th}}$. Obtaining $\Delta m^2 \sim \mathcal{O}(10^{-4} \text{ eV}^2)$ requires, for $m_\nu \sim 1 \text{ eV}$ the correction $I_{e\mu(\tau)}^{\text{th}} \sim 10^{-(4-5)}$ which is possible in the MSSM and gives $r \sim 10^{-(1-2)}$ i.e. acceptable s_{13} .

$\mathcal{O}(\Delta m_{ab}/\max(|m_{\nu_3}|, |m_\nu|))$, i.e. small if the hierarchy of neutrino masses is large. If only flavour diagonal threshold corrections are present the solar mass squared splitting is given by²¹

$$\Delta m_{21}^2 \approx 4m_\nu^2 \left[(U_{32}^2 - U_{31}^2) \tilde{I}_\tau + (U_{12}^2 - U_{11}^2) \tilde{I}_e \right]. \quad (5.14)$$

The interesting aspect of this situation is that the (generically) dominant first order contribution to Δm^2 proportional to $\tilde{I}_\tau = \tilde{I}_\tau^{\text{rg}} + \tilde{I}_\tau^{\text{th}}$ vanishes for $U_{32}^2 = U_{31}^2$, i.e. for $s_{13} = \pm(s_{23}/c_{23})(c_{12} \pm s_{12})/(c_{12} \mp s_{12})$, a special case of which is the bimaximal mixing solution with $s_{13} = 0$ (also the second term in eq. (5.14) then vanishes). This has been discussed in refs. [50, 49, 36] in connection with the possibility of realizing the VO solution to the solar neutrino problem, within the inversely hierarchical pattern $m_{\nu_3}^2 \ll m_\nu^2 \sim 3 \times 10^{-3} \text{ eV}^2$. Note that for generic mixing angles the RG corrections would always give too large a Δm^2 .

Since the second order correction ($\propto \tilde{I}_\tau^2$) to Δm^2 is proportional to m_{ν_3} , the right Δm^2 for the VO solution can be obtained by appropriately tuning $m_{\nu_3}^2 \gtrsim 0$ and/or due to the threshold effects.

As discussed in [36], with $m_{\nu_1} = -m_{\nu_2}$, $m_{\nu_3}^2 \ll m_{\nu_1}^2 \sim 3 \times 10^{-3} \text{ eV}^2$, the correct Δm^2 for the LAMSW (or SAMSW) solutions can be obtained from the RG running. From eq. (5.14) we get

$$\Delta m^2 \approx 4m_\nu^2 \left| s_{23}^2 \cos 2\theta_{12} + s_{13} \sin 2\theta_{23} \sin 2\theta_{12} \right| \tilde{I}_\tau^{\text{rg}} \quad (5.15)$$

and for the LAMSW solution, with $|s_{13}| \sim 0.1$, $\Delta m^2 \sim \mathcal{O}(10^{-5}) \text{ eV}^2$ can be obtained for $\tan \beta \gtrsim 30$ ($\tan \beta \gtrsim 10$ for the SAMSW solution). For smaller values of $\tan \beta$ one has to investigate potentially larger threshold corrections, which give [60]

$$\begin{aligned} \Delta m^2 &\approx 4m_\nu^2 \left| \sin 2\theta_{12} (c_{23} I_{e\mu}^{\text{th}} - s_{23} I_{e\tau}^{\text{th}}) \right| && \text{LAMSW} \\ \Delta m^2 &\approx 4m_\nu^2 \left| \cos 2\theta_{12} (\tilde{I}_e^{\text{th}} - \frac{1}{2} \sin 2\theta_{23} I_{\mu\tau}^{\text{th}}) \right| && \text{SAMSW.} \end{aligned} \quad (5.16)$$

For $\Delta m^2 \sim \mathcal{O}(10^{-5}) \text{ eV}^2$ one needs therefore corrections $\tilde{I}^{\text{th}} \sim 10^{-3} - 10^{-2}$ which is definitely too big a value for $I_{e\mu}^{\text{th}}$ in the MSSM and also impossible for $I_{e\tau}^{\text{th}}$ and $I_{\mu\tau}^{\text{th}}$ (at least within the validity of the mass insertion approximation, it would require $|\delta_{LL}^{e(\mu)\tau}| > 1$).

Our discussion here does not cover fully the potential role of quantum corrections for the two-fold degeneracy pattern. Several other possibilities do exist, depending on the chosen solar solution and we refer the reader to the literature [61, 60, 56] for further details.

²¹We choose to work with $\tilde{I}_A \equiv I_A - I_\mu$.

6 Conclusions

For every theory of neutrino masses, for a meaningful comparison with experimental information, it is necessary to discuss quantum corrections. This is relatively easy if the effective low energy theory is the SM or its supersymmetric extension and the neutrino masses enter via the effective operator (3.11). In Sections 4 and 5 we reviewed the formalism for including quantum correction in those cases. Next we applied that formalism to several different classes of the neutrino mass sector at the scale M , hypothetically given by the theory of neutrino masses. They split into two broad groups. The first one is characterized by small quantum corrections that may eventually be important for precision tests of the future theory. However, they are irrelevant at the level of present, qualitative considerations. Here belongs the hierarchical pattern and the partly degenerate pattern with opposite CP parities of the (almost) degenerate neutrinos. For those mass matrices, quantum corrections cannot substantially alter the structures present at the scale M , so the agreement with experimental data has to be assured by the boundary values at M . Equal masses $m_{\nu_1} = -m_{\nu_2}$, at the scale M are possible. Quantum corrections can explain the observed Δm^2 .

The other group consists of partly degenerate pattern with the same CP parities of the (almost) degenerate neutrinos and of the (approximately) degenerate structures. Large quantum corrections can originate either from the RG evolution or from low energy threshold corrections (in the MSSM). They never change qualitatively the mass eigenvalue pattern, although they may explain their observed splitting. However, large quantum corrections always lead to a “fixed point” relation for the mixing angles. The sufficient condition is that for at least one pair of neutrinos $(m_{\nu_a} + m_{\nu_b})/(m_{\nu_a} - m_{\nu_b}) \gg 1$. It is interesting that the transition from small (qualitatively irrelevant) quantum corrections to the “fixed point” behaviour is very abrupt. So, to a good approximation there are those two and only two physical situations.

A fixed point relation is always one equation for three angles (if CP is conserved) and makes their low energy values dependent on only two boundary conditions for the angles at the scale M . If the dominant quantum corrections come from the RG evolution (i.e. originate from the large τ Yukawa coupling) or also from several configurations of the low energy threshold corrections, the “fixed point” relation links small θ_{13} angle (constrained by the CHOOZ experiment) to a small angle responsible for the mixing of solar neutrinos. If presently favoured bimaximal mixing was confirmed by future experimental data, all mass patterns leading to large RG corrections and the regions of the MSSM parameter space leading to the same “fixed point” relation would be ruled out.

There are, however, other “fixed point” relations, generated by some special sfermion

mass configurations and/or by flavour non-conserving effects in the slepton sector, that are consistent with bimaximal mixing and small θ_{13} angle. Such solutions are phenomenologically interesting as, at the same time, they explain the observed mass squared differences, as the effect of quantum corrections, with degenerate spectrum at the scale M .²² However, the simple Ansatz at the scale M needs a deeper theoretical justification.

Quantum corrections do not explain the origin of the neutrino masses and do not replace its theory. Nevertheless, they are important piece of the overall picture. They will constrain strongly the acceptable mass structures once the experimental ambiguities are resolved.

Acknowledgments

The work of P.H.Ch. was supported partially by the EC Contract HPRN-CT-2000-00148 for years 2000-2004 and by the Polish State Committee for Scientific Research grant 2 P03B 060 18 for years 2000-2001. The work of S.P. was supported partially by the EC Contract HPRN-CT-2000-00152 for years 2000-2004 and by the Polish State Committee for Scientific Research grant 5 P03B 119 20 for years 2001-2002. P.H.Ch. Would also like to thank the CERN Theory group for hospitality during the writing of this article.

Appendix A

In Section 3, instead of the familiar Dirac bispinors we used the two-component Weyl spinor notation. It is particularly convenient for dealing with Majorana particles. Here we explain this notation shortly. More details can be found in modern textbooks on QFT (see e.g. [16, 64]).

In four dimensions, the Lorentz group or more precisely its covering group $SL(2, C)$ has two non-equivalent complex two-dimensional representations denoted as $(1/2, 0)$ and $(0, 1/2)$. The Grassmann fields (or fermionic field operators) transforming according to these representations are conventionally written as λ_α and $\bar{\chi}^{\dot{\alpha}}$ and called²³ left- and right-handed spinors, respectively. Since the complex conjugation of a left-handed spinor λ_α

²²It has also been discussed in the literature [52, 53] that quantum corrections generated in the full theory above the scale M , can break the degeneracy already at the scale M and lead to correct mass squared differences and mixing angles at the electroweak scale. For an interesting link between neutrino masses and flavour changing processes see ref. [63].

²³Somewhat incorrectly; properly they should be called left- and right-chiral.

transforms as a right-handed one $((\lambda_\alpha)^* \sim \bar{\lambda}_{\dot{\alpha}})$, the fermion content of any Lagrangian can be specified by listing only the left-handed spinors used for its construction.

If the two left-handed fields λ_α and χ_β transform as representations R and R^* , respectively under the final unbroken symmetry group (global or local) of the theory, they can be combined to form a Dirac bispinor:

$$\psi_{(\lambda)} = \begin{pmatrix} \lambda_\alpha \\ \bar{\chi}^{\dot{\beta}} \end{pmatrix}, \quad \bar{\psi}_{(\lambda)} = (\chi^\alpha, \bar{\lambda}^{\dot{\beta}}), \quad (\text{A.1})$$

transforming as R and R^* , respectively. The raising and lowering of Weyl spinor indices is done with the help of the antisymmetric tensors $\epsilon^{\alpha\beta}$, $\epsilon_{\alpha\beta}$, $\epsilon_{\dot{\alpha}\dot{\beta}}$ and $\epsilon^{\dot{\alpha}\dot{\beta}}$:

$$\lambda^\alpha = \epsilon^{\alpha\beta} \lambda_\beta, \quad \lambda_\alpha = \lambda^\beta \epsilon_{\beta\alpha}, \quad \bar{\chi}_{\dot{\alpha}} = \epsilon_{\dot{\alpha}\dot{\beta}} \bar{\chi}^{\dot{\beta}}, \quad \bar{\chi}^{\dot{\alpha}} = \bar{\chi}_{\dot{\beta}} \epsilon^{\dot{\beta}\dot{\alpha}}. \quad (\text{A.2})$$

The two kinetic terms for λ and χ can be then rewritten in the familiar form

$$\mathcal{L}_{\text{kin}} = i\bar{\lambda}\bar{\sigma}^\mu\partial_\mu\lambda + i\bar{\chi}\bar{\sigma}^\mu\partial_\mu\chi = i\bar{\lambda}\bar{\sigma}^\mu\partial_\mu\lambda + i\chi\sigma^\mu\partial_\mu\bar{\chi} + (\text{total der}) = i\bar{\psi}_{(\lambda)}\gamma^\mu\partial_\mu\psi_{(\lambda)} \quad (\text{A.3})$$

where the Dirac matrices γ^μ in the Weyl representation are constructed as

$$\gamma^\mu = \begin{pmatrix} 0 & \sigma^\mu_{\alpha\dot{\beta}} \\ \bar{\sigma}^{\mu\dot{\alpha}\beta} & 0 \end{pmatrix} \quad (\text{A.4})$$

with $\sigma^\mu \equiv (I, \boldsymbol{\sigma})$, $\bar{\sigma}^\mu \equiv (I, -\boldsymbol{\sigma})$ ($\boldsymbol{\sigma}$'s are the Pauli matrices). For such a pair of Weyl fields also a Dirac mass term can be constructed

$$\mathcal{L}_{\text{mass}} = -m(\lambda^\alpha\chi_\alpha + m\bar{\lambda}_{\dot{\alpha}}\bar{\chi}^{\dot{\alpha}}) = -m\bar{\psi}_{(\lambda)}\psi_{(\lambda)} \quad (\text{A.5})$$

If the field λ (χ) has no left-handed partner transforming in the complex conjugate representation R^* (R), it is convenient to introduce chiral Dirac bispinors

$$\psi_{(\lambda)L} = \begin{pmatrix} \lambda_\alpha \\ 0 \end{pmatrix}, \quad \psi_{(\chi)R} = \begin{pmatrix} 0 \\ \bar{\chi}^{\dot{\alpha}} \end{pmatrix}. \quad (\text{A.6})$$

For chiral Dirac bispinors e.g. $\psi_{(\lambda)L} = P_L\psi_{(\lambda)L}$ etc., where $P_L \equiv (1 - \gamma^5)/2$. Note that (see eq.(A.1)),

$$\overline{\psi_{(\lambda)L}} = (0 \quad \bar{\lambda}_{\dot{\alpha}}), \quad \overline{\psi_{(\chi)R}} = (\chi^\alpha \quad 0). \quad (\text{A.7})$$

The typical Yukawa coupling of a scalar field ϕ in the representation R_ϕ and two left-handed Weyl spinors λ and χ transforming as representations R_λ and R_χ , respectively (such that $1 \subset R_\phi \times R_\lambda \times R_\chi$) can be written as (we omit the Clebsch-Gordan coefficients)

$$\begin{aligned} \mathcal{L}_{\text{Yuk}} &= -Y\phi\lambda\chi - Y^*\phi^\dagger\bar{\lambda}\bar{\chi} \\ &= -Y\phi\overline{\psi_{(\chi)R}}\psi_{(\lambda)L} - Y^*\phi^\dagger\overline{\psi_{(\lambda)L}}\psi_{(\chi)R} \end{aligned} \quad (\text{A.8})$$

The Yukawa part of the SM Lagrangian (3.1) and the SM mass terms (3.3) are the example of (A.8) and (A.5), respectively, with fields u^c , d^c and e^c playing the role of χ , and q and l (or u , d and e) playing the role of λ .

Finally, Weyl spinor fields λ_α which are singlets of all unbroken symmetries of the theory can form 4-component Majorana bispinors

$$\psi_{(\lambda)\text{Maj}} = \begin{pmatrix} \lambda^\alpha \\ \bar{\lambda}^{\dot{\beta}} \end{pmatrix}. \quad (\text{A.9})$$

Of course in this case $\psi_{(\lambda)} = C\bar{\psi}_{(\lambda)}^T \equiv \psi_{(\lambda)}^c$ which means that the field is self-conjugate. One also has (up to raising or lowering indices)

$$\lambda = \psi_{(\lambda)L} = \overline{\psi_{(\lambda)R}} \quad \text{and} \quad \bar{\lambda} = \psi_{(\lambda)R} = \overline{\psi_{(\lambda)L}}. \quad (\text{A.10})$$

For such a field a Majorana mass term can be formed

$$\begin{aligned} \mathcal{L}_{\text{Maj}} &= -\frac{1}{2}m(\lambda\lambda + \bar{\lambda}\bar{\lambda}) = -\frac{1}{2}m\left(\overline{\psi_{(\lambda)R}}\psi_{(\lambda)L} + \overline{\psi_{(\lambda)L}}\psi_{(\lambda)R}\right) \\ &\equiv -\frac{1}{2}m\bar{\psi}_{(\lambda)}\psi_{(\lambda)} \equiv -\frac{1}{2}m\psi_{(\lambda)}^T C\psi_{(\lambda)} \end{aligned} \quad (\text{A.11})$$

where C is the charge conjugation matrix. The Majorana mass term (3.8) is precisely of this form with ν^c playing the role of λ and the Yukawa coupling (3.7) is usually written as

$$\Delta\mathcal{L}_{\text{Yuk}} = -\epsilon_{ij}H_i\overline{\psi_{(\nu^c)R}}\mathbf{Y}_\nu\psi_{(l)L} - \epsilon_{ij}H_i^*\overline{\psi_{(\lambda)L}}\mathbf{Y}_\nu^\dagger\psi_{(\nu^c)R} \quad (\text{A.12})$$

Appendix B

In this Appendix we recall the well known RGEs for Yukawa coupling matrices defined in eq. (3.1) in the SM and in the MSSM. In the SM they read [25, 24]:

$$\begin{aligned} \frac{d}{dt}\mathbf{Y}_u &= \mathbf{Y}_u \left[-8g_3^2 - \frac{9}{4}g_2^2 - \frac{17}{12}g_Y^2 + T + \frac{3}{2}(\mathbf{Y}_u^\dagger\mathbf{Y}_u - \mathbf{Y}_d^\dagger\mathbf{Y}_d) \right], \\ \frac{d}{dt}\mathbf{Y}_d &= \mathbf{Y}_d \left[-8g_3^2 - \frac{9}{4}g_2^2 - \frac{5}{12}g_Y^2 + T + \frac{3}{2}(\mathbf{Y}_d^\dagger\mathbf{Y}_d - \mathbf{Y}_u^\dagger\mathbf{Y}_u) \right], \\ \frac{d}{dt}\mathbf{Y}_e &= \mathbf{Y}_e \left[-\frac{9}{4}g_2^2 - \frac{15}{4}g_Y^2 + T + \frac{3}{2}\mathbf{Y}_e^\dagger\mathbf{Y}_e \right], \end{aligned} \quad (\text{B.1})$$

where

$$t \equiv \frac{1}{16\pi^2} \ln \left(\frac{Q}{M_Z} \right), \quad (\text{B.2})$$

$$T \equiv \text{Tr} \left(3\mathbf{Y}_u^\dagger\mathbf{Y}_u + 3\mathbf{Y}_d^\dagger\mathbf{Y}_d + \mathbf{Y}_e^\dagger\mathbf{Y}_e \right) \quad (\text{B.3})$$

and the gauge couplings g_3 , g_2 and g_Y evolve according to

$$\frac{d}{dt}g_i = b_i g_i^3 \quad i = 3, 2, Y \quad (\text{B.4})$$

with $b_3 = -7$, $b_2 = -19/6$ and $b_Y = 41/6$.

In the MSSM one finds instead:

$$\begin{aligned} \frac{d}{dt}\mathbf{Y}_u &= \mathbf{Y}_u \left[-\frac{16}{3}g_3^2 - 3g_2^2 - \frac{13}{9}g_Y^2 + \text{Tr} \left(3\mathbf{Y}_u^\dagger \mathbf{Y}_u \right) + 3\mathbf{Y}_u^\dagger \mathbf{Y}_u + \mathbf{Y}_d^\dagger \mathbf{Y}_d \right], \\ \frac{d}{dt}\mathbf{Y}_d &= \mathbf{Y}_d \left[-\frac{16}{3}g_3^2 - 3g_2^2 - \frac{7}{9}g_Y^2 + \text{Tr} \left(3\mathbf{Y}_d^\dagger \mathbf{Y}_d + \mathbf{Y}_e^\dagger \mathbf{Y}_e \right) + 3\mathbf{Y}_d^\dagger \mathbf{Y}_d + \mathbf{Y}_u^\dagger \mathbf{Y}_u \right], \\ \frac{d}{dt}\mathbf{Y}_e &= \mathbf{Y}_e \left[-3g_2^2 - 3g_Y^2 + \text{Tr} \left(3\mathbf{Y}_d^\dagger \mathbf{Y}_d + \mathbf{Y}_e^\dagger \mathbf{Y}_e \right) + 3\mathbf{Y}_e^\dagger \mathbf{Y}_e \right], \end{aligned} \quad (\text{B.5})$$

and the factors b_i change to $b_3 = -3$, $b_2 = +1$ and $b_Y = 11$.

For completeness we give here also the RGEs above the scale M i.e. for the theory whose set of fermion fields includes additional three $SU_L(2) \times U_Y(1)$ singlet neutrino fields ν^{cA} ($A = 1, 2, 3$) and whose Lagrangian is identical to the one for the effective theory valid below M except for the Yukawa interaction (3.7) and the Majorana mass terms (3.8).

If the theory above the M scale extends the SM, then T given in eq. (B.3) has to be replaced by [65, 39, 35]

$$T \equiv \text{Tr} \left(3\mathbf{Y}_u^\dagger \mathbf{Y}_u + 3\mathbf{Y}_d^\dagger \mathbf{Y}_d + \mathbf{Y}_e^\dagger \mathbf{Y}_e + \mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu \right), \quad (\text{B.6})$$

the last equation in (B.1) should be replaced by

$$\frac{d}{dt}\mathbf{Y}_e = \mathbf{Y}_e \left[-\frac{9}{4}g_2^2 - \frac{15}{4}g_Y^2 + T + \frac{3}{2} \left(\mathbf{Y}_e^\dagger \mathbf{Y}_e - \mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu \right) \right] \quad (\text{B.7})$$

and the neutrino Yukawa matrix RGE reads:

$$\frac{d}{dt}\mathbf{Y}_\nu = \mathbf{Y}_\nu \left[-\frac{9}{4}g_2^2 - \frac{3}{4}g_Y^2 + T - \frac{3}{2} \left(\mathbf{Y}_e^\dagger \mathbf{Y}_e - \mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu \right) \right]. \quad (\text{B.8})$$

In addition, the Majorana mass matrix also runs [52]:

$$\frac{d}{dt}\mathbf{M}_{\text{Maj}}^{KL} = \mathbf{M}_{\text{Maj}}^{KJ} \left(\mathbf{Y}_\nu \mathbf{Y}_\nu^\dagger \right)^{LJ} + \left(\mathbf{Y}_\nu \mathbf{Y}_\nu^\dagger \right)^{KJ} \mathbf{M}_{\text{Maj}}^{JL}. \quad (\text{B.9})$$

If the low energy theory is the MSSM, then [39, 35] in the first equation of (B.5)

$$\text{Tr} \left(3\mathbf{Y}_u^\dagger \mathbf{Y}_u \right) \rightarrow \text{Tr} \left(3\mathbf{Y}_u^\dagger \mathbf{Y}_u + \mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu \right), \quad (\text{B.10})$$

the last equation of (B.5) is replaced by

$$\frac{d}{dt}\mathbf{Y}_e = \mathbf{Y}_e \left[-3g_2^2 - 3g_Y^2 + \text{Tr} \left(3\mathbf{Y}_d^\dagger \mathbf{Y}_d + \mathbf{Y}_e^\dagger \mathbf{Y}_e \right) + 3\mathbf{Y}_e^\dagger \mathbf{Y}_e + \mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu \right] \quad (\text{B.11})$$

and the neutrino Yukawa matrix RGE reads:

$$\frac{d}{dt}\mathbf{Y}_\nu = \mathbf{Y}_\nu \left[-3g_2^2 - g_Y^2 + \text{Tr} \left(3\mathbf{Y}_u^\dagger \mathbf{Y}_u + \mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu \right) + 3\mathbf{Y}_e^\dagger \mathbf{Y}_e + \mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu \right]. \quad (\text{B.12})$$

Finally, the Majorana mass matrix running is dictated by [53]:

$$\frac{d}{dt}\mathbf{M}_{\text{Maj}}^{KL} = 2\mathbf{M}_{\text{Maj}}^{KJ} \left(\mathbf{Y}_\nu \mathbf{Y}_\nu^\dagger \right)^{LJ} + 2 \left(\mathbf{Y}_\nu \mathbf{Y}_\nu^\dagger \right)^{KJ} \mathbf{M}_{\text{Maj}}^{JL}. \quad (\text{B.13})$$

References

- [1] K. Nishikawa, plenary talk at the *International EPS Conference on High Energy Physics*, Budapest, Hungary, July 2001.
- [2] G. Altarelli and F. Feruglio, *Phys. Lett.* **B439** (1998), 112, *JHEP* **9811:021** (1998), *Phys. Rept.* **320** (1999), 295; G. Altarelli, F. Feruglio and I. Masina, *Phys. Lett.* **B472** (2000), 382.
- [3] The Particle Data Group, *Eur. Phys. J.* **C15** (2000), 1.
- [4] C. Giunti, C.W. Kim, J.A. Lee and U.W. Lee, *Phys. Rev.* **D48** (1993), 4310; J. Rich, *Phys. Rev.* **D48** (1993), 4318; C. Cardall *Phys. Rev.* **D61** (2000), 073006; M. Zralek, *Acta Phys. Pol.* **B29** (1998), 3925.
- [5] S.M. Bilenkii, C. Giunti and W. Grimus, *Prog. Part. Nucl. Phys.* **43** (1999), 1.
- [6] S.P. Mikheyev and A.Yu. Smirnov, *Sov. J. Nucl. Phys.* **42** (1985), 913; L.F. Wolfenstein, *Phys. Rev.* **D17** (1978), 2369.
- [7] Y. Fukuda et al. (The Superkamiokande Collaboration), *Phys. Rev. Lett.* **81** (1998), 1562.
- [8] S. Fukuda et al. (The Superkamiokande Collaboration), *Phys. Rev. Lett.* **85** (2000), 3999.
- [9] C. Bemporad, G. Gratta and P. Vogel, hep-ph/0107277.
- [10] C. Athanassopoulos et al. (The LSND Collaboration), *Phys. Rev. Lett.* **77** (1996), 3082, **81** (1998), 1774.
- [11] Q.R. Ahmad et al. (The SNO Collaboration), *Phys. Rev. Lett.* **87** (2001), 071301.

- [12] S. Fukuda et al. (The Superkamiokande Collaboration), *Phys. Rev. Lett.* **86** (2001), 5651, 5656.
- [13] M. Apollonio et al. (The CHOOZ Collaboration), *Phys. Lett.* **B466** (1999), 415.
- [14] Z. Maki, M. Nakagawa and S. Sakata, *Prog. Theor. Phys.* **28** (1962), 870.
- [15] L. Baudis et al. (Heidelberg-Moscow Collaboration), *Phys. Rev. Lett.* **83** (1999), 41;
A. Dietz (Heidelberg-Moscow Collaboration), talk presented at the *3rd International Conference on Dark Matter in Astro and Particle Physics*, Heidelberg, Germany, Jul 2000 (hep-ph/0103062).
- [16] S. Pokorski, *Gauge Field Theories*, 2nd edition, Cambridge University Press, Cambridge, 2000, Chapters 1 and 12.
- [17] M. Gell-Mann, P. Ramond and R. Slansky, in proceedings of the *Supergravity Stony Brook Workshop*, New York, 1979, eds. P. Van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam); T. Yanagida, in proceedings of the *Workshop on Unified Theories and Baryon Number in the Universe*, Tsukuba, 1979, eds. A. Sawada and A. Sugamoto, KEK Report No. 79-18.
- [18] R.N. Mohapatra, preprint UMDPP-02-002, 2001 (hep-ph/0107264).
- [19] S. Weinberg, *Phys. Rev. Lett.* **43** (1979), 1566.
- [20] K.S. Babu and C.N. Leung, preprint OSU-HEP-01-02, UDHEP-02-01, hep-ph/0106054.
- [21] R. Hempfling, *Nucl. Phys.* **B478** (1996), 3; E. Nardi, *Phys. Rev.* **D55** (1997), 5772; E.J. Chun, S.K. Kang, C.W. Kim and U.W. Lee, *Nucl. Phys.* **B544** (1999), 89; E.J. Chun and S.K. Kang *Phys. Rev.* **D61** (2000), 075012.
- [22] T.P. Cheng, E. Eichten and L.-F. Li, *Phys. Rev.* **D9** (1974) 2259.
- [23] M.E. Machaček and M.T. Vaughn *Nucl. Phys.* **B236** (1984), 221.
- [24] V. Barger, M.S. Berger and P. Ohmann, *Phys. Rev.* **D47** (1993), 1093.
- [25] K.S. Babu, *Z. Phys.* **C35** (1987), 69.
- [26] M. Olechowski and S. Pokorski, *Phys. Lett.* **231B** (1989) 165; *ibid.* **257B** (1991), 388.

- [27] V. Barger, M.S. Berger and P. Ohmann, *Phys. Rev.* **D47** (1993), 2038.
- [28] L.F. Wolfenstein *Phys. Rev. Lett.* **51** (1983), 1945.
- [29] S. Antusch, M. Drees, J. Kersten, M. Lindner and M. Ratz, preprint TUM-HEP-424/01 (hep-ph/0108005).
- [30] C. Wetterich, *Nucl. Phys.* **B187** (1981), 343; A. Ioannisian *Sov. J. Nucl. Phys.* **51** (1990), 511.
- [31] P.H. Chankowski and Z. Pluciennik, *Phys. Lett.* **B316** (1993), 312.
- [32] K.S. Babu, C.N. Leung and J. Pantaleone, *Phys. Lett.* **B319** (1993), 191.
- [33] P.H. Chankowski, *Phys. Rev.* **D41** (1990), 2877.
- [34] J. Ellis and S. Lola, *Phys. Lett.* **B458** (1999), 310.
- [35] N. Haba, Y. Matsui, N. Okamura and M. Sugiura, *Eur. Phys. J.* **C10** (1999), 677.
- [36] P.H. Chankowski, W. Królikowski and S. Pokorski, *Phys. Lett.* **B473** (2000), 109.
- [37] J.A. Casas, J.R. Espinosa, A. Ibarra and I. Navarro, *Nucl. Phys.* **B573** (2000), 652.
- [38] M. Tanimoto, *Phys. Lett.* **B360** (1995), 41
- [39] N. Haba, N. Okamura and M. Sugiura, *Prog. Theor. Phys.* **103** (2000), 367.
- [40] K.R.S. Balaji, A. Dighe, R.N. Mohapatra and M.K. Parida, *Phys. Rev. Lett.* **84** (2000), 5034.
- [41] K.R.S. Balaji, R.N. Mohapatra, M.K. Parida and E.A. Paschos, *Phys. Rev.* **D63**:113002 (2001).
- [42] J. Ellis, G.K. Leontaris, S. Lola and D.V. Nanopoulos, *Eur. Phys. J.* **C9** (1999) 389.
- [43] S. Lola, proceedings of the 6th Hellenic School and Workshop on Elementary Particle Physics, Corfu, Greece, September 1998 (hep-ph/9903203).
- [44] M. Carena, J. Ellis, S. Lola, C.E.M. Wagner, *Eur. Phys. J.* **C12** (2000), 507.
- [45] E. Ma, *J. Phys.* **G25** (1999), L97-L100.
- [46] N. Haba, Y. Matsui and N. Okamura, *Prog. Theor. Phys.* **103** (2000), 807; T.K. Kuo, J. Pantaleone and G.-H. Wu.

- [47] T.K. Kuo, J. Pantaleone and G.-H. Wu, hep-ph/0103131.
- [48] N. Haba, Y. Matsui, N. Okamura and M. Sugiura, *Prog. Theor. Phys.* **103** (2000), 145.
- [49] R. Barbieri, G.G. Ross and A. Strumia, *JHEP* **9910:020** (1999).
- [50] J.A. Casas, J.R. Espinosa, A. Ibarra and I. Navarro, *JHEP* **9909:015** (1999).
- [51] A.S. Dighe and A.S. Joshipura, preprint CERN-TH-2000-301 (hep-ph/0010079).
- [52] J.A. Casas, J.R. Espinosa, A. Ibarra and I. Navarro, *Nucl. Phys.* **B556** (1999), 3.
- [53] J.A. Casas, J.R. Espinosa, A. Ibarra and I. Navarro, *Nucl. Phys.* **B569** (2000), 82.
- [54] N. Haba and N. Okamura, *Eur. Phys. J.* **C14** (2000), 347.
- [55] K.R.S. Balaji, A. Dighe, R.N. Mohapatra and M.K. Parida, *Phys. Lett.* **B481** (2000), 33.
- [56] P.H. Chankowski and P. Wasowicz, preprint IFT-01/28, (hep-ph/0110237).
- [57] L.I. Schiff, *Quantum Mechanics*, 3rd edition, McGraw-Hill Book Company, New York, 1968.
- [58] P.H. Chankowski, A. Ioannisian, S. Pokorski and J.W.F. Valle, *Phys. Rev. Lett.* **86** (2001), 3488.
- [59] E. Dudas, S. Pokorski and C.A. Savoy, *Phys. Lett.* **B369** (1996), 255; E. Dudas, Ch. Grojean, S. Pokorski and C.A. Savoy *Nucl. Phys.* **B481** (1996), 85; A.G. Cohen, D.B. Kaplan and A.E. Nelson, *Phys. Lett.* **B388** (1996), 588; Ph. Brax and C.A. Savoy, *J. High. Energy Phys.* **0007:048** (2000).
- [60] E. J. Chun, *Phys. Lett.* **B505** (2001) 155.
- [61] E.J. Chun and S. Pokorski, *Phys. Rev.* **D62:053001** (2000).
- [62] F. Gabbiani, E. Gabrieli, A. Masiero and L. Silvestrini, *Nucl. Phys.* **B447** (1996), 321; see also, M. Misiak, S. Pokorski and J. Rosiek, in *Heavy Flavours II*, eds. A.J. Buras and M. Lindner, World Scientific Publishing Co., Singapore 1998.
- [63] S. Lavignac, I. Masina and C.A. Savoy, preprint SACLAY-T01/066 (hep-ph/0106245).

- [64] M.E. Peskin and D.V. Schroeder, *An Introduction to Quantum Field Theory*, Addison Wesley, Reading Massachussets 1997.
- [65] B. Grzadkowski and M. Lindner, *Phys. Lett.* **B193** (1987), 71; Y.F. Pirogov and O. Zenin, hep-ph/9808396.